

SULFUR IN THE EARLY MARTIAN ATMOSPHERE REVISITED: EXPERIMENTS WITH A 3-D GLOBAL CLIMATE MODEL. L. Kerber¹, F. Forget¹, R. Wordsworth². ¹Laboratoire de Météorologie Dynamique du CNRS, Université Paris 6, Paris, France (Kerber@lmd.jussieu.fr), ²Department of Geosciences, University of Chicago, Chicago, Illinois 60637

Introduction: Data returned from the surface of Mars during the 1970s revealed intriguing geological evidence for a warmer and wetter early martian climate. Dendritic valley networks were discovered by Mariner 9 on ancient Noachian terrain [1], indicating that liquid water had flowed across the surface in the distant past. Since this time, geological investigations into early Martian history have attempted to ascertain the nature and level of activity of the early Martian hydrological cycle [e.g. 2-5] while atmospheric modeling efforts have focused on how the atmosphere could be warmed to temperatures great enough to sustain such activity [see 6-7 for reviews].

Geological and spectroscopic investigations have refined the history and chronology of Noachian Mars over time, and circulation of liquid water has been invoked to explain several spatially and temporally distinct morphological and chemical signatures found in the geological record. Detections of iron and magnesium-rich clays are widespread in the oldest Martian terrains, suggesting a period of pH-neutral aqueous alteration [e.g., 8]. Valley network incision also took place during the Noachian period [9]. Some chains of river valleys and craters lakes extend for thousands of kilometers, suggesting temperatures at least clement enough for sustained ice-covered flow [3, 10]. The commencement of valley network incision is not well constrained, but the period of Mg/Fe clay formation appears to have ended before the termination of valley network formation, as the visible fluvial systems appear to have remobilized existing clays rather than forming them [5,8]. There is also evidence that the cessation of valley network formation was abrupt [11]. Towards the end of the Noachian, erosion rates appear to have been significantly higher than during subsequent periods, a process that has also been attributed to aqueous processes [12]. A period of sulfate formation followed, likely characterized by acidic, evaporitic playa environments [8].

A successful working model for the early Martian atmosphere and hydrosphere must be able not only to produce conditions suitable for liquid water at the surface, but also to explain how the nature of this aqueous activity changed over time and eventually diminished. There are two major end-member hypotheses: first, that early Mars was wet and warm, with a sustained greenhouse that made it possible for liquid water to be stable on the surface for extended periods [e.g., 2, 12-14], and second, that early Mars was generally cold, and that most of the aqueous alteration took place underground [3,5] or during transient warm periods tied to impact cratering [15], or volcanism [16]. In both of these scenarios it is generally agreed that in order to make valley networks and sulfate deposits, a hydrological cycle is needed which is able to recycle water from the lowlands back to the highlands (i.e., the one-time emptying of a regional aquifer would not be sufficient to create the observed features) [4,17]. This would require some precipitation to fall on the southern highlands, either flowing overland or filtering into groundwater aquifers.

In both cases, volcanic gases (especially SO₂) have been suggested as a possible way of creating either a sustained or transient greenhouse. Several researchers have tested the addition of SO₂ to climate models in order to assess whether it would provide an adequate amount of greenhouse warming to allow liquid water to flow across the surface [18-21], with differing results. Postawko and Kuhn [18] found a warming effect of 14 K in a 0.1 bar atmosphere with an SO₂ abundance of 1000 ppm. A total greenhouse warming effect of 44 K was found using the same mixing ratio in a much thicker atmosphere (1 bar), but the amount of SO₂ needed to create this same mixing ratio in a thicker atmosphere was deemed unreasonable, and it was concluded that SO₂ could not be responsible for any appreciable early greenhouse. Johnson et al. [20] used a 3-D global circulation model and found a warming of 15-25 K for 245 ppm of SO₂ in a dry 0.5 bar atmosphere. Tian et al. [21] used a 1-D model to explore a wide range of SO₂ mixing values and CO₂ partial pressures, finding a warming of around ~25 K for 100 ppm in a 0.5 bar atmosphere with a fully saturated troposphere (~40 K for a 1 bar atmosphere). These authors also included the effect of sulfate aerosol particles, which caused a dramatic cooling effect which more than canceled the warming caused by the SO₂ gas [21].

Here we reconsider the efficacy of a sulfur-induced greenhouse in early Noachian history using the LMD (Laboratoire de Météorologie Dynamique) 3-D Generic Climate Model (LMD-GCM), exploring the effects of SO₂, H₂S, and sulfate aerosols on the surface temperature.

Methods: The GCM used in this study was developed for general planetary applications, and includes generalized radiative transfer and cloud physics [e.g., 22]. The model was adapted specifically to early Mars conditions and used by Forget et al. [7] to explore the effects of a CO₂ atmosphere from 0.1 to 7 bars, including the effects of different obliquities, orbital parameters, cloud microphysical parameters, atmospheric dust loading, and surface properties. A companion paper exploring the effects of a full water cycle including surface-atmosphere interactions and water ice clouds [23] concluded that even with the greenhouse warming provided by water vapor and water-ice clouds, above-freezing conditions were not met for sustained periods. However, this study found that over the long term, ice tended to be transported from the lowlands to the highlands, providing a pathway for the recharge of water. If SO₂ warming was responsible for valley network formation, we might expect to find high temperatures where valley networks are found, to cause preferential melting in these areas. If SO₂ warming was responsible for allowing ground-water-fed playa lakes to exist in a liquid form long enough to evaporate [4], we might expect higher temperatures in areas where sulfates were observed (e.g., Arabia Terra).

Simulations were run at 0.5 bars with a 32x32x15 grid, corresponding to a resolution of 11.25° longitude by 5.625° latitude in the horizontal with 15 vertical levels from the surface to ~50 km. Other resolutions and pressures were also

explored to test sensitivity. The correlated-k method was used to produce a database of coefficients for use by the radiative transfer code, using 80 spectral bands in the longwave and 36 in the shortwave. Real and imaginary indices for H_2SO_4 were taken from the HITRAN database, using the values from [24] for wavelengths between 0.2 and 0.337 μm and between 25 μm and 50 μm , while results from [25] were used for wavelengths between 0.360 μm and 25 μm (following [21] to allow for comparison). Simulations with H_2O were run with a relative humidity profile derived from the full, self-consistent water cycle (including CO_2 and water ice clouds) used in [23]. The relative humidity in this profile is 45% near the surface with the hygropause near 8 km.

Results: Simulations were run in a dry CO_2 atmosphere with various SO_2 and H_2S mixing ratios: 0 ppm, 10 ppm, 100 ppm, 1000 ppm, and 10000 ppm (Fig. 1).




Figure 1.

Even in the case where a large amount of SO_2 is added to the atmosphere, the annual average surface temperature does not rise above freezing anywhere on the planet (local maximum temperatures do exceed freezing). H_2S provides significantly less warming than SO_2 . Figure 2 shows maps of average annual surface temperature for different scenarios: CO_2 only, $\text{H}_2\text{O}+\text{CO}_2$, CO_2+SO_2 , and $\text{CO}_2+\text{H}_2\text{O}+\text{SO}_2$. The addition of water vapor greatly increases the greenhouse effect of the SO_2 , though global average annual temperatures are still below freezing (245 K). The warmest areas do not coincide with the locations of the valley networks, but there is warming in Arabia Terra, which could have aided in the formation of sulfate deposits.

Additional simulations were run in dry CO_2 atmospheres with well-mixed H_2SO_4 0.5 μm particles with opacities from $\tau=0.2$ to 10, producing cooling from a few degrees to ~ 23 K for $\tau=10$. A single layer of H_2SO_4 aerosols of the same opacity produced similar results. Larger particles produced less

cooling with very large particles (10 μm) producing mild warming. The greenhouse warming produced by our model is therefore significantly less than that produced by that of Johnson et al. [20], but slightly greater (5-10 K) than that of Tian et al. [21]. The cooling produced by sulfate particles is significantly less than that of Tian et al. [21]. While a sustained greenhouse is not created, local and seasonal melting would be facilitated provided that the SO_2 lifetime was sufficient.

References: [1] Sharp, R.P., Malin, M.C. (1975) GSA Bull. 86, 593-609. [2] Baker, V.R. et al. (1991) Nature 352, 589-594. [3] Squyres, S.W., Kasting, J.F. (1994) Science 265, 744-749. [4] Andrews-Hanna, J. et al., 2010 JGR 115, E06002. [5] Ehlmann, B. et al. (2011) Nature 479, 53-60. [6] Haberle, R.M. (1998) JGR 103, 28,467-28,479. [7] Forget, F. et al. (2013) Icarus 222, 81-99. [8] Bibring, J-P. et al., Science 312, 400-404. [9] Carr, M.H. (1996) *Water on Mars*. Oxford Univ. Press, 229pp. [10] Fassett, C.I., Head, J.W. (2008) Icarus 198, 37-56. [11] Fassett, C.I., Head, J.W. (2008) Icarus 195, 61-89. [12] Hynek, B.M., Phillips, R.J. (2001) Geology 29, 407-410. [13] Craddock, R.A., Howard, A.D. (2002) JGR 107, 5111. [14] Halevy, I. et al. (2007) Science 318, 1903-1907. [15] Segura, T.L., et al. (2002) Science 298, 1977-1988. [16] Halevy, I., and Head, J.W. (2012) LPSC 43, Abs. 1908. [17] Goldspiel, J.M., Squyres, S.W. (1991) Icarus 89, 392-401. [18] Postawko, S.E., Kuhn, W.R. (1986) LPSC Proc. 91, D431-D438. [19] Mischna, M.A. et al. (2011) AGU Fall Meet. Abs. P11G-04. [20] Johnson, S.S., et al. (2008) JGR 113, E08005. [21] Tian, F., et al. (2010) EPSL 295, 412-418. [22] Wordsworth, R et al. (2011) Astrophys. J. Lett. 733:L48. [23] Wordsworth, R., et al. (2013) Icarus 222, 1-19. [24] Hummel, J.R., et al. (1988) AFGL-TR-88-0166, Hanscom AFB. [25] Palmer, K. F., Williams, D. (1975) Applied Optics 14, 208-219.




Figure 2. Average annual surface temperature, 0.5 bars.