

THE EARLIEST STAGES OF HIGH MASS STAR FORMATION – METHANOL MASER INSIGHTS

V. Minier¹, P. André¹, F. Motte¹, N. Peretto¹, R.S. Booth², J.E. Conway², M.R. Pestalozzi³, M.G. Burton⁴, T. Hill⁴, S.N. Longmore⁴, C.R. Purcell⁴, A.J. Walsh⁴, R. Cesaroni⁵, F. Herpin⁶, J.M. de Buizer⁷ and M. Elitzur⁸

¹Service d'Astrophysique, CEA Saclay, 91191 Gif-sur-Yvette, France – Vincent.Minier@cea.fr

²Onsala Space Observatory, 439 92 Onsala, Sweden

³School of Physics, Astronomy and Maths, University of Hertfordshire, Hatfield, UK

⁴School of Physics, University of New South Wales, Sydney 2052, NSW, Australia

⁵Osservatorio Astrofisico di Arcetri, Largo E. Fermi 5, 50125 Firenze, Italy

⁶Observatoire de Bordeaux, Floirac, France

⁷GEMINI Observatory, Casilla 603, La Serena, Chile

⁸Department of Physics and Astronomy, University of Kentucky, USA

We report on recent multi-scale observational studies of the earliest stages of high mass ($>8 M_{\odot}$) star formation using methanol MASERs as astronomical probes. Methanol masers can provide unique information about densely populated, embedded protoclusters in which precursors of O and B stars form.

Tracers of high mass star-forming complexes in the Galactic plane: The brightest methanol masers are detected in radio frequencies at 6.7 and 12.2 GHz and are not affected by extinction in high mass star-forming regions. They were originally detected in the environment of very active star-forming complexes of HII regions a decade ago [1]. Since their discovery, more than 500 methanol maser sites have been located in the Galactic plane through searches toward IRAS colour selected sources [2], OH and H₂O masers [3] as well as through unbiased full-sampling surveys [4,5]. Interestingly, 6.7-GHz methanol masers have not been detected toward low mass star-forming regions despite very sensitive searches [6]. All these results strongly suggest that 6.7-GHz methanol masers are exclusively associated with high mass star-forming complexes in the Galactic plane [5] (Fig. 1a).

Tracers of massive protoclusters: Several complementary studies have been undertaken to identify the nature of the relationship between the maser sites and the star-forming regions. Methanol masers arise from deeply embedded (>10 mag), massive ($>50 M_{\odot}$), cold (20-50 K) and luminous ($>10^3 L_{\odot}$) molecular clumps (~ 0.5 pc in diameter) [7,8,9]. Various classes of object can be identified among these molecular clumps based on their thermal dust emission in submillimetre and infrared. Methanol masers are associated with both mid-infrared dark and bright clumps, a combination often seen in a unique complex (e.g. Fig. 1b). The mid-IR dark clumps are characterised by a Spectral Energy Distribution of cold dust emission that peaks at longer wavelengths (~ 20 -30 K) than that of bright IR clumps (~ 40 -50 K). Mid-IR dark clumps might then represent early stages of the clustered star formation process. Many bright IR clumps are radio

quiet in terms of free-free continuum emission. These results indicate that within high mass star-forming complexes methanol masers trace massive protoclusters of young stellar objects in earlier evolutionary phases than in HII regions.

Tracers of high mass protostars: High angular resolution observations have demonstrated that methanol masers are in fact generally not associated with strong radio continuum emission from ultra-compact HII regions [10]. However, methanol maser sites appear to coincide with hot molecular cores as well as with hyper compact and very weak radio sources in many cases [11]. They are also detected in mid-IR at sub-arcsec resolution and pinpoint bright objects with luminosities ($\sim 10^4 L_{\odot}$), radius (~ 100 AU) and masses ($\sim 10 M_{\odot}$) consistent with those expected for high mass young stellar objects [12,13] (e.g. Fig. 1c,d). VLBI observations of methanol masers have revealed that they often form lines with velocity gradients along them [14] which are interpreted as protostellar Keplerian disks seen edge-on [15] (e.g. Fig. 1e). Alignments with outflows are also observed [13,14]. These results strongly suggest that CH₃OH masers trace precursors of UC HII regions, i.e. sites of massive protostars.

References:

- [1] Menten K. (1991) ApJ, 380, L75
- [2] Walsh A.J., Hyland A. R., Robinson G., Burton M.G. (1997) MNRAS, 291, 261
- [3] Caswell J.L., Vaile R.A., Ellingsen S.P., Whiteoak J.B., Norris R.P. (1995) MNRAS, 272, 96
- [4] Ellingsen S.P., von Bibra M.L., McCulloch P.M. et al. (1996) MNRAS, 280, 378
- [5] Pestalozzi M.R., Minier V., Booth R.S. (2005) A&A, 432, 737
- [6] Minier V., Ellingsen S., Norris R.P., Booth R.S., (2003) A&A, 403, 1095
- [7] Minier V., Burton M.G., Hill T. et al. (2005) A&A, 429, 945
- [8] Hill T., Burton M.G., Minier V. et al. (2005) MNRAS, in press
- [9] Purcell C.R. et al. (2005) MNRAS, submitted

[10] Walsh A.J., Burton M.G., Hyland A.R., Robinson G. (1998) MNRAS, 301, 640
 [11] Minier V., Conway J.E., Booth R.S. (2001) A&A, 369, 278
 [12] Longmore S.N. et al. (2005) MNRAS, submitted

[13] De Buizer J.M., Minier V. (2005) ApJ, 628, 151
 [14] Minier V., Booth R.S., Conway J.E. (2000) A&A, 362, 1093
 [15] Pestalozzi M.R., Elitzur M., Conway, J.E., Booth R.S. (2004) ApJ, 603, 113




Figure 1: **a.** Distribution of methanol maser sites in the Galactic plane overlaid on the CO(1-0) map of molecular clouds. **b.** NGC 7538 – a complex of high mass star formation. Methanol masers (white crosses) are associated with the bright IR source IRS 1 in NGC 7538, as well as with the deeply embedded IRS 9 and the dark cloud NGC 7538 S. **c.** Close-up composite JHK image with 2MASS of IRS1-3. The contours represent 18- μ m thermal dust emission detected with GEMINI/MICHELLE. **d.** Close-up of the mid-IR image. Black contours are radio continuum emission at 22 GHz. Blue symbols are methanol masers. **e.** Close-up of the maser line. Colours vary with velocities.