

10057

Ilmenite Basalt (high K)

919 grams



Figure 1: Photo of whole rock 10057,0 during preliminary examination. NASA photo # S69-46287. Sample is about 11 cm across.

Introduction

All of the Apollo 11 basalts have high TiO_2 and low SiO_2 contents. Apollo 11 did not sample a very wide area and it appears that only 2 lava flows may have been sampled, distinguished by higher K, and slightly higher REE, contents in one flow compared with the other. There is also a correlation with cosmic ray exposure age.

10057 belongs to the group of samples known as “high K, Apollo 11 basalts”. Although they range in texture, they are remarkable similar in composition and are probably from the same basalt flow (Beaty and Albee 1978). The range of texture is attributed to differences in cooling rate. 10057 is fine-grained and cooled rather quickly (Grove and Beaty 1980). The crystallization age is about 3.6 b.y. and the cosmic ray exposure age is about 54 m.y.

Petrography

Lunar basalt 10057 is composed of intergrown subhedral to anhedral plagioclase, euhedral to subhedral ilmenite, subhedral pyroxene and a residual fine-grained mesostasis composed of plagioclase, cristobalite and potassic glass. 10057 has a high abundance (~10%) of vesicles (figure 1). LSPET (1969) termed this group of vesicular, fine-grained rock “type A basalts”. James and Wright (1972) termed these rocks “intersertal ilmenite basalts”. Schmitt et al. (1970) termed 10057 as a “fine-grained, vesicular to vuggy, granular basalt.” Grove and Beaty (1980) term them “antiophitic”.




Figure 2: Photomicrograph of thin section of 10057 (plane polarized light). NASA #76-26315.

Mineralogy

Pyroxene: Pyroxenes in 10057 are chemically zoned in major and minor elements (Lovering and Ware 1970, Essene et al. 1970 and Beatty and Albee 1978) (figure 3). Large grains have pigeonite cores with augite rims trending towards iron rich, but without the extreme iron enrichment of other lunar basalts. Essene et al. (1970) studied the variation of Ti and Al as a function of Mg/Fe in pyroxene in 10057.

Plagioclase: The plagioclase in 10057 ranges in composition from An₇₄ to An₈₁ (Reid et al. 1970, Beatty

and Albee 1978). Steward et al. (1970) obtain An₇₅ from cell size measurements.

Opacues: Ilmenite in 10057 is about the same size (40 microns) as pyroxene or plagioclase (Lovering and Ware 1970).

Chemistry

Lunar basalt 10057 was widely distributed and analyzed by numerous persons (tables 1, 2). Figures 4

Mineralogical Mode for 10057

	Haggerty et al. 1970	Beatty and Albee 1978
Olivine		<0.04
Pyroxene	50.9 vol. %	50.8
Plagioclase	19.2	24
Opacues	15.7	15.5
Cristobalite		1.05
Troilite		0.42
Phosphate		0.21
Mesostasis	3.3	8.04
Metal		0.04
Vesicles	11	
Armalcolite		absent




Figure 3: Olivine and pyroxene composition diagram for 10057 (data replotted from Lovering and Ware 1970 and Beaty and Albee 1977).

and 5 show that 10057 is a typical high-K, high-Ti Apollo 11 basalt.

Radiogenic age dating

Papanastassiou et al. (1970) dated 10057 as 3.63 ± 0.02 b.y. by the Rb-Sr internal isochron method. Davis et al. (1971) and Tatsumoto (1970) found that they could not date 10057 by the $^{39}\text{Ar}/^{40}\text{Ar}$ or U/Pb methods.

Cosmogenic isotopes and exposure ages

O'Kelley et al. (1970) determined the cosmic ray induced activity of ^{22}Na (41 dpm/kg), ^{26}Al (75 dpm/kg), ^{46}Sc (10 dpm/kg), ^{54}Mn (32 dpm/kg) and ^{56}Co (31 dpm/kg). Perkins et al. (1970) found ^{22}Na (43 dpm/kg), ^{26}Al (84 dpm/kg), ^{46}Sc (11 dpm/kg), ^{54}Mn (41 dpm/kg) and ^{56}Co (19 dpm/kg). Wrigley and Quaide (1970) measured significant ^{22}Na (49 dpm/kg) and ^{26}Al (75 dpm/kg) in 10057 due to recent cosmic ray events.

The cosmic ray exposure ages for 10057 were reviewed by Hintenberger et al. (1971) and Srinivasan (1974). Hohenberg et al. (1970) determined 34 ± 5 m.y. by ^{83}Kr , Marti (1970) and Marti and Lugmair (1971) determined 47 ± 2 m.y. by ^{83}Kr , Bochsler et al. (1971) reported 52.5 m.y. Hintenberger et al. (1971) and Guggisberg et al. (1979) determined 58 m.y. and 52 m.y. by ^{38}Ar . Arvidson et al. (1975) reported a ^{81}Kr exposure age of 52.5 m.y. (determined by Schwaller 1971). Eugster et al. (1984) determined 54.2 m.y. Srinivasan (1974) used the Xe data from Marti et al.

Summary of Age Data for 10057

	Rb-Sr	Ar-Ar	K-Ar
Papanastassiou et al. 1970	3.63 ± 0.02 b.y.		

Caution: Age not corrected for "new" Rb decay constant.




Figure 4: Composition of lunar basalts showing position for 10057 (data from table 1).




Figure 5: Normalized rare-earth-element diagram for high-K basalt 10057 (data from Wanke et al. 1970 and Haskin et al. 1970 only).

(1970) and Hohenberg et al. (1970) to calculate 34 and 54 m.y. respectively.

Other Studies

The rare gas contents and isotopic ratios of 10057 were reported by Hohenberg et al. (1970), Hintenberger et al. (1971), Bogard et al. (1971) and Eugster et al. (1984).

Oxygen isotope ratios were reported for whole rock and mineral separates of 10057 by O'Neil and Adams (1970) and Taylor and Epstein (1970).

Table 1a. Chemical composition of 10057.

reference weight	Engel 70	Wanke 70	Wasson 70	Wrigley 70	Ganapathy 70	Anders 71	Morrison 70	Perkins 70	O'Kelly 70
SiO ₂ %	39.79	(a) 40.43					46	(d)	
TiO ₂	11.44	(a) 10.84	(b)				7.56	(b)	
Al ₂ O ₃	10.84	(a) 7.56	(b)				10.84	(b)	
FeO	19.35	(a) 18.01	(b)				20.2	(b)	
MnO	0.2	(a) 0.23	(b)				0.22	(b)	
MgO	7.65	(a) 6.96	(b)				6.13	(b)	
CaO	10.08	(a) 11.75	(b)				14.13	(b)	
Na ₂ O	0.54	(a) 0.4	(b)				0.47	(b)	
K ₂ O	0.32	(a) 0.24	(b)				0.26	(b)	0.28 (e) 0.31 (e)
P ₂ O ₅	0.17	(a)					0.09	(c)	
S %									
sum									
Sc ppm	100	(f) 87	(b)				84	(b)	
V	66	(f)					40	(b)	
Cr	2400	(f) 2160	(b)				2100	(b)	
Co	21	(f) 25.4	(b)		27.2	(b) 30	(b) 24	(b)	
Ni	7	(f) <10	(b)				40	(b)	
Cu	11	(f) 4.3	(b)		3.52	(b)	5.5	(b)	
Zn					1.7	(b) 1.75	(b) 2.9	(b)	
Ga		5.2	(b) 4.9	(b)			4.7		
Ge ppb			< 70	(b)					
As									
Se						0.181	(b)		
Rb		5.2	(b)		3.68	(b)	4.8	(b)	
Sr	130	(f) 100	(b)				130	(b)	
Y	180	(f)					210	(c)	
Zr	400	(f)			560	(b)	360	(b)	
Nb							42	(c)	
Mo							0.4	(b)	
Ru									
Rh									
Pd ppb		10	(b)		7.3	(b)			
Ag ppb					0.69	(b)			
Cd ppb					3.15	(b) 3.5	(b)		
In ppb		2.7	(b) 3		3.2	(b)			
Sn ppb									
Sb ppb									
Te ppb					8	(b)			
Cs ppm		0.2	(b)		0.159	(b)	0.2	(b)	
Ba	130	(f) 208	(b)				280	(b)	
La		25	(b)				31	(b)	
Ce		79	(b)				83	(b)	
Pr		9	(b)				22	(c)	
Nd		60	(b)				66	(b)	
Sm		20	(b)				24	(b)	
Eu		1.8	(b)				2.1	(b)	
Gd		30	(b)				26	(b)	
Tb		5	(b)				5.6	(b)	
Dy		24	(b)				42	(c)	
Ho		5.5	(b)				8	(b)	
Er		16	(b)				32	(c)	
Tm							2.3	(b)	
Yb	16	(f) 16.8	(b)				26	(b)	
Lu		2.15	(b)				2.2	(b)	
Hf		16.9	(b)				15	(b)	
Ta		2	(b)				1.2	(b)	
W ppb		430	(b)				0.42	(b)	
Re ppb									
Os ppb									
Ir ppb			0.1		0.023	(b) 0.009	(b)		
Pt ppb									
Au ppb		1.6	< 0.3		0.017	(b) 0.013	(b)		
Th ppm		3.94	(b)	3.27	(e)		4.5	(b) 3.6	(e) 3.3 (e)
U ppm		0.8	(b)	0.97	(e)		0.56	(b) 0.95	(e) 0.79 (e)

technique (a) wet chem. (b) INAA, RNAA, (c) SSMS, (d) AA, (e) radiation counting, (f) emission spec.

Table 1b. Chemical composition of 10057.

<i>reference weight</i>	Smales71	Turekian70	Kharkar71	Annell 70	Tatsumoto70 Haskin 70 Baedecker70	LSPET 70	Duncan76	Neal2001	
SiO ₂ %	41.4					36	(c) 40.67	(g)	
TiO ₂	10.9	8.2	12.18	(b)		12.5	(c) 11.35	(g)	
Al ₂ O ₃	8.1					11	(c) 7.99	(g)	
FeO	19.1		18.65	(b)		20	(c) 19.4	(g)	
MnO	0.23	0.25	0.23	(b) 0.28	(c)	0.49	(c) 0.233	(g)	
MgO	7.4					9.5	(c) 7.65	(g)	
CaO	10.4		9.37	(b)		10	(c) 10.75	(g)	
Na ₂ O	0.54	0.55	0.55	(b)		0.54	(c) 0.57	(g)	
K ₂ O	0.33	0.43				0.18	(c) 0.287	(g)	
P ₂ O ₅							0.197	(g)	
S %							0.26	(g)	
<i>sum</i>									
Sc ppm	86		90	(b) 99	(c)	110	(c)	77.2	(h)
V				65	(c)	50	(c) 47	(g) 45.5	(h)
Cr	2250		2290	(b) 2790	(c)	6500	(c) 2374	(g) 2063	(h)
Co			26	(b) 30	(c)	22	(c) 25	(g) 26.8	(h)
Ni				6.1	(c)	25	(c) <2	(g) 5.91	(h)
Cu				5.7	(c)			60.5	(h)
Zn								75.6	(h)
Ga				5	(c) 4.9	(b)		4.19	(h)
Ge ppb					<0.2	(b)			
As									
Se		0.12							
Rb				4.7	(c)	6	(c) 6.5	(g) 5.27	(h)
Sr				140	(c)	230	(c) 166	(g) 144	(h)
Y				165	(c)	310	(c) 170	(g) 176	(h)
Zr				635	(c)		517	(g) 523	(h)
Nb				29	(c)		28.7	(g) 26.7	(h)
Mo		0.1						0.07	(h)
Ru									
Rh									
Pd ppb									
Ag ppb		52							
Cd ppb									
In ppb									
Sn ppb									
Sb ppb								20	(h)
Te ppb									
Cs ppm								0.36	(h)
Ba				440	(c)	180	(c) 319	(g) 261	(h)
La		25	23	(b) 26	(c) 28.2	(b)		23.4	(h)
Ce			69.6	(b)	75	(b)		83.3	(h)
Pr								11.1	(h)
Nd					69	(b)		59.4	(h)
Sm		17	14.3	(b)	20.8	(b)		20.1	(h)
Eu		2.5	2.4	(b)	2.18	(b)		1.95	(h)
Gd					26	(b)		26.4	(h)
Tb					5	(b)		4.53	(h)
Dy		37	29.9	(b)	34.7	(b)		29.3	(h)
Ho								6.5	(h)
Er					19	(b)		17.8	(h)
Tm								2.43	(h)
Yb			13.2	(b)	18.8	(b) 6	(c)	17	(h)
Lu			2.48	(b)	2.66	(b)		2.26	(h)
Hf			18.1	(b)				13.6	(h)
Ta			1.7	(b)				1.76	(h)
W ppb								520	(h)
Re ppb									
Os ppb									
Ir ppb									
Pt ppb									
Au ppb		6.4							
Th ppm					3.415	(a)		2.77	(h)
U ppm		0.47			0.865	(a)		0.76	(h)

technique (a) IDMS, (b) INAA, (c) emission spec., (g) XRF, (h) ICP-MS

Table 2. Light and/or volatile elements for 10057

	Engel 70	Wanke 70	Annell 70	Morrison 70	Ganapathy 70
Li ppm	14	14	17	8	
Be			3.3	2.5	
B		0.8		4	
C					
S					
F ppm				70	
Cl		12		50	
Br ppb				100	25.2
I					
Pb ppm					
Hg ppb					
Tl ppb					1.09
Bi ppb					0.27



Figure 6: Space Window at US National Cathedral in Washington DC with piece of 10057 located in center of rose window.




Figure 7: Photo of thin slice of 10057 located in Space Window.




Figure 8: Photo of 10057 after sawing into chunks. NASA S75-33923. Cube is 1 cm.

Housley et al. (1970) determined the Mössbauer spectra.

Processing

Apollo 11 samples were originally described and cataloged in 1969 and “recataloged” by Kramer et al. (1977). There are 12 thin sections.

A piece of 10057 (large area thin section) is in the center of the Space Window of the National Cathedral (figures 6 and 7). In 1976 the sample was sawn at various odd angles into several chunks (figures 8 and 9).

Table 2	U ppm	Th ppm	K ppm	Rb ppm	Sr ppm	Nd ppm	Sm ppm	technique
Perkins 1970	0.95	3.6	2300					rad. Cout.
O'Kelley 1970	0.79	3.3	2550					rad. Cout.
Haskin 1970						69	20.8	





Figure 9: Group photo of pieces of 10057. NASA S75-20522. Cube is 1 cm.



References for 10057

- Arvidson R., Crozaz G., Drozd R.J., Hohenberg C.M. and Morgan C.J. (1975) Cosmic ray exposure ages of features and events at the Apollo landing sites. *The Moon* **13**, 259-276.
- Anders E., Ganapathy R., Keays R.R., Laul J.C. and Morgan J.W. (1971) Volatile and siderophile elements in lunar rocks: Comparison with terrestrial and meteoritic basalts. *Proc. 2nd Lunar Sci. Conf.* 1021-1036.
- Anders E., Ganapathy R., Krahenbuhl U. and Morgan J.W. (1973) Meteoritic material on the Moon. *The Moon* **8**, 3-24.
- Annell C.S. and Helz A.W. (1970) Emission spectrographic determination of trace elements in lunar samples from Apollo 11. *Proc. Apollo 11 Lunar Sci. Conf.* 991-994.
- Baedecker P.A., Schaudy R., Elzie J.L., Kimberlin J., and Wasson J.T. (1971) Trace element studies of rocks and soils from Oceanus Procellarum and Mare Tranquilitatis. *Proc. 2nd Lunar Sci. Conf.* 1037-1061.
- Beaty D.W. and Albee A.L. (1978) Comparative petrology and possible genetic relations among the Apollo 11 basalts. *Proc. 9th Lunar Planet. Sci. Conf.* 359-463.
- Bochsler P., Eberhardt P., Geiss J., Loosli H., Oeschger H. and Wahlen M. (1971) Tritium in lunar material. *Proc. Second Lunar Sci. Conf.* 1803-1812.
- Bogard D.D., Funkhouser J.G., Schaeffer O.A. and Zahringer J. (1971) Noble gas abundances in lunar material-cosmic ray spallation products and radiation ages from the Sea of Tranquillity and the Ocean of Storms. *J. Geophys. Res.* **76**, 2757-2779.
- Davis P.K., Lewis R.S. and Reynolds J.H. (1971) Stepwise heating analysis of rare gases from pile-irradiated rocks 10044 and 10057. *Proc. Second Lunar Sci. Conf.* 1693-1703.
- Engel A.E.J. and Engel Celeste G. (1970) Lunar rock compositions and some interpretations. *Proc. Apollo 11 Lunar Sci. Conf.* 1081-1084.
- Engel A.E.J., Engel C.G., Sutton A.L. and Myers A.T. (1971) Composition of five Apollo 11 and Apollo 12 rocks and one Apollo 11 soil and some petrogenetic considerations. *Proc. 2nd Lunar Sci. Conf.* 439-448.
- Essene E.J., Ringwood A.E. and Ware N.G. (1970) Petrology of the lunar rocks from Apollo 11 landing site. *Proc. Apollo 11 Lunar Sci. Conf.* 385-397.
- Eugster O., Eberhardt P., Geiss J., Grögler N. and Schwaller H. (1984b) Cosmic ray exposure histories and ²³⁵U-¹³⁶Xe dating of Apollo 11, Apollo 12, and Apollo 17 mare basalts. *Proc. 15th Lunar Planet. Sci. Conf.* in *J. Geophys. Res.* **89**, C171-C181.
- Ganapathy R., Keays R.R., Laul J.C. and Anders E. (1970) Trace elements in Apollo 11 lunar rocks: Implications for meteorite influx and origin of moon. *Proc. Apollo 11 Lunar Sci. Conf.* 1117-1142.
- Grove T.L. and Beatty D.W. (1980) Classification, experimental petrology and possible volcanic histories of the Apollo 11 high-K basalts. *Proc. 11th Lunar Planet. Sci. Conf.* 149-177.
- Guggisberg S., Eberhardt P., Geiss J., Grogler N., Stettler A., Brown G.M. and Peckett A. (1979) Classification of the Apollo-11 basalts according to Ar³⁹-Ar⁴⁰ ages and petrological properties. *Proc. 10th Lunar Planet. Sci. Conf.* 1-39.
- Haggerty S.E., Boyd F.R., Bell P.M., Finger L.W. and Bryan W.B. (1970) Opaque minerals and olivine in lavas and breccias from Mare Tranquilitatis. *Proc. Apollo 11 Lunar Sci. Conf.* 513-538.
- Haskin L.A., Allen R.O., Helmke P.A., Paster T.P., Anderson M.R., Korotev R.L. and Zweifel K.A. (1970) Rare earths and other trace elements in Apollo 11 lunar samples. *Proc. Apollo 11 Lunar Sci. Conf.* 1213-1231.
- Hintenberger H., Weber H.W. and Takaoka N. (1971) Concentrations and isotopic abundances of the rare gases in lunar matter. *Proc. 2nd Lunar Sci. Conf.* 1607-1625.
- Hohenberg C.M., Davis P.K., Kaiser W.A., Lewis R.S. and Reynolds J.H. (1970) Trapped and cosmogenic rare gases from stepwise heating of Apollo 11 samples. *Proc. Apollo 11 Lunar Sci. Conf.* 1283-1309.

- Housley R.M., Blander M., Abdel-Gawad M., Grant R.W. and Muir A.H. (1970) Mossbauer spectroscopy of Apollo 11 samples. *Proc. Apollo 11 Lunar Sci. Conf.* 2251-2268.
- Hurley P.M. and Pinson W.H. (1971) Whole-rock Rb-Sr isotopic age relationships in Apollo 11 lunar samples. *Proc. Apollo 11 Lunar Sample Conference* 1311-1315.
- James O.B. and Jackson E.D. (1970) Petrology of the Apollo 11 ilmenite basalts. *J. Geophys. Res.* **75**, 5793-5824.
- James O.B. and Wright T.L. (1972) Apollo 11 and 12 mare basalts and gabbros: Classification, compositional variations and possible petrogenetic relations. *Geol. Soc. Am. Bull.* **83**, 2357-2382.
- Kharkar D.P. and Turekian K.K. (1970) Neutron activation analysis of milligram quantities of Apollo 11 lunar rocks and soil. *Proc. Apollo 11 Lunar Sci. Conf.* 1659-1664.
- Kharkar D.P. and Turekian K.K. (1971) Analyses of Apollo 11 and Apollo 12 rocks and soils by neutron activation. *Proc. 2nd Lunar Sci. Conf.* 1301-1305.
- Kramer F.E., Twedell D.B. and Walton W.J.A. (1977) **Apollo 11 Lunar Sample Information Catalogue** (revised). Curator's Office, JSC 12522
- Lovering J.F. and Ware N.G. (1970a) Electron probe microanalyses of minerals and glasses in Apollo 11 lunar samples. *Proc. Apollo 11 Lunar Sci. Conf.* 633-654.
- LSPET (1969) Preliminary examination of lunar samples from Apollo 11. *Science* **165**, 1211-1227.
- Marti K., Lugmair G.W. and Urey H.C. (1970) Solar wind gases, cosmic ray spallation products and the irradiation history of Apollo 11 samples. *Proc. Apollo 11 Lunar Sci. Conf.* 1357-1367.
- Marti K. and Lugmair G.W. (1971) Kr^{81} -Kr and Kr - Ar^{40} ages, cosmic-ray spallation products and neutron effects in lunar samples from Oceanus Procellarum. *Proc. 2nd Lunar Sci. Conf.* 1591-1605.
- Morrison G.H., Gerard J.T., Kashuba A.T., Gangadharam E.V., Rothenberg A.M., Potter N.M. and Miller G.B. (1970) Elemental abundances of lunar soil and rocks. *Proc. Apollo 11 Lunar Sci. Conf.* 1383-1392.
- Neal C.R. (2001) Interior of the moon: The presence of garnet in the primitive deep lunar mantle. *J. Geophys. Res.* **106**, 27865-27885.
- O'Kelley G.D., Eldridge J.S., Schonfeld E. and Bell P.R. (1970) Primordial radionuclide abundances, solar proton and cosmic ray effects and ages of Apollo 11 lunar samples by non-destructive gamma-ray spectrometry. *Proc. Apollo 11 Lunar Sci. Conf.* 1407-1424.
- O'Neil J.R. and Adams L.H. (1970) Oxygen isotope analyses of selected Apollo 11 materials. *Proc. Apollo 11 Lunar Sci. Conf.* 1425-1427.
- Papanastassiou D.A., Wasserburg G.J. and Burnett D.S. (1970a) Rb-Sr ages of lunar rocks from the Sea of Tranquillity. *Earth Planet. Sci. Lett.* **8**, 1
- Papanastassiou D.A. and Wasserburg G.J. (1971a) Lunar chronology and evolution from Rb-Sr studies of Apollo 11 and 12 samples. *Earth Planet. Sci. Lett.* **11**, 37-62.
- Perkins R.W., Rancitelli L.A., Cooper J.A., Kaye J.H. and Wogman N.A. (1970) Cosmogenic and primordial radionuclide measurements in Apollo 11 lunar samples by nondestructive analysis. *Proc. Apollo 11 Lunar Sci. Conf.* 1455-1469.
- Reid A.M., Frazer J.Z., Fujita H. and Everson J.E. (1970a) Apollo 11 samples: Major mineral chemistry. *Proc. Apollo 11 Lunar Sci. Conf.* 749-761.
- Reid A.M., Frazer J.Z., Fujita H. and Everson J.E. (1970b) Chemical composition of the major phases in Apollo 11 lunar samples. SIO ref. 70-4 Univ. Calif. San Diego
- Schmitt H.H., Lofgren G., Swann G.A. and Simmons G. (1970) The Apollo 11 samples: Introduction. *Proc. Apollo 11 Lunar Science Conf.* 1-54.
- Schwaller H., Eberhardt P., Geiss J., Graf H. and Grogler N. (1971) The $^{78}Kr/^{83}Kr - ^{131}Xe/^{126}Xe$ correlation in Apollo 12 rocks. *Earth Planet. Sci. Lett.* **12**, 167-169.
- Smales A.A., Mapper D., Webb M.S.W., Webster R.K., Wilson J.D., and Hislop J.S. (1971) Elemental composition of lunar surface material (part 2). *Proc. Seond Lunar Sci. Conf.* 1253-1258.

Stewart D.B., Appleman D.E., Huebner J.S. and Clark J.R. (1970) Crystallography of some lunar plagioclase. *Science* **167**, 634-635.

Srinivasan B. (1974) Lunar breccia 14066: $^{81-83}\text{Kr}$ exposure age, evidence for fissionogenic xenon from ^{244}Pu and rate of production of spallogenic ^{126}Xe . *Proc. 5th Lunar Sci. Conf.* 2033-2044.

Sutton R.L. and Schaber G.G. (1971) Lunar locations and orientations of rock samples from Apollo missions 11 and 12. *Proc. 2nd Lunar Sci. Conf.* 17-26.

Tatsumoto M. (1970) Age of the Moon: An isotopic study of U-Th-Pb systematics of Apollo 11 lunar samples - II. *Proc. Apollo 11 Lunar Sci. Conf.* 1595-1612.

Taylor H.P. and Epstein S. (1970a) O18/O16 ratios of Apollo 11 lunar rocks and minerals. *Proc. Apollo 11 Lunar Sci. Conf.* 1613-1626.

Wänke H., Rieder R., Baddenhausen H., Spettler B., Teschke F., Quijano-Rico M. and Balacescu A. (1970) Major and trace elements in lunar material. *Proc. Apollo 11 Lunar Sci. Conf.* 1719-1727.

Wasson J.T. and Baedeker P.A. (1970) Ga, Ge, In, Ir, and Au in lunar terrestrial and meteoritic basalts. *Proc. Apollo 11 Lunar Sci. Conf.* 1741-1750.

Wrigley R.C. and Quaide W.L. (1970) Al26 and Na22 in lunar surface materials: Implications for depth distribution studies. *Proc. Apollo 11 Lunar Sci. Conf.* 1751-1755.