

14313
Regolith Breccia
 144 grams



Figure 1: Photo of 14313. Sample is 6 cm across. NASA S71-29131.




Figure 2: Photo of 14313 before it was collected (AS14-68-9466).

CDR Man, that pile of rocks – beautiful, right to your left. Oh, just the right size. Don't walk over them!
 LMP No, I'm trying to stay away from them. Yes __ it's bigger than we thought. Al, we'll grab sample that one; I'll get you another one here.
 CDR Okay, bag 27 Nancy. (#14313)
 LMP And another documented sample – a larger documented sample than we thought we were getting here, Fredo. Again, it was a burried rock: and it's too big for the sample bag; so, it'll go into the weigh bag. (#14301)

LMP We're approaching Triplet from the east, that's North Triplet from the east. There's a little rock field down here – a small boulder field, Al, to get a documented sample from.




Figure 3: Map of Apollo 14 with location of 14313.





Figure 4: Thin section photomicrograph of 14313 showing seriate texture. Scale 0.5 cm. NASA S71-25455.



*Figure 5: Photomicrograph of thin section 14314,9 showing basalt clast. Scale = 2.8 mm across.
Photo by C Meyer.*

Introduction

14313 was collected at station G1 on the rim of North Triplet Crater about 150 meters from the LM (figure 3). Like nearby sample 14301, it was also partially “buried” (figure 2), but has been shown to have had a complex exposure history of micrometeorite bombardment indicating “tumbling” while on the regolith surface (Morrison et al. 1972).

14313 is a fine-grained regolith breccia with seriate grain size distribution. It has few large clasts and is mostly matrix. The Apollo 14 regolith breccias (vitric matrix breccias) are slightly more aluminous than the the Fra Mauro breccias (crystalline matrix breccias). They also have intermediate values of C, N, ^4He and other elements from the ancient solar wind indicating that they contain a soil component.

Petrography

Lunar sample 14313 is a coherent polymict breccia that has had a complex history of “comminution and re-agglomeration” (Floran et al. 1972). The matrix is composed primarily of fine particles of brown glass (figure 4). Clasts include (1) noritic rock fragments, (2) monomineralic fragments, (3) microbreccia clasts, (4) glassy fragments including chondrules and %) rare mare basalt clasts (figure 9). There is evidence of breccia-in-breccia relations indicating multiple generations of breccia formation.

Drozd et al. (1976) reported a high percentage of agglutinates in 14313, but Simon et al. (1989) found considerably less. Other indications of maturity are relatively high C, N and ^4He contents. Fruland (1983) and Simon et al. (1981) classified 14313 as a regolith breccia.




Figure 6: Composition of Apollo 14 breccias with 14313.




Figure 7: Normalized rare-earth-element diagrams for 14313 and for KREEP.

King et al. (1972) studied the chondrules in 14313. Delano (1987), Simon et al. (1989) and Wentworth and McKay (1991) reported numerous glass analyses.

Significant Clast

Micronorite (or basalt)

Floran et al. (1972) described a basaltic clast in 14313 as a “micronorite” (figure 9).

Alkali Anorthosite? ,70

Warren et al. (1983) reported a clast of alkali anorthosite.

Chemistry

Boynton et al. (1975), Helmke et al. (1972), Simon et al. (1989) and Hughes et al. (1973) reported the composition of 14313 (table 1). 14313 is relatively high in Al (figure 6) and very high in REE (figure 7).

Moore et al. (1972) reported 130 and 170 ppm carbon for two subsamples of 14301 (figure 10). Becker and Clayton (1975) reported 400×10^{-4} ppm He and 47 ppm nitrogen (indications of a substantial solar wind component).

Radiogenic age dating

Alexandra and Kahl (1974) could not obtain an “age” for breccia 14313 by the Ar-Ar technique (figure 11), but state that it is assumed to be 3.7 ± 0.2 b.y.

Other Studies

Becker R.H. and Clayton R.N. (1975) determined the He (400×10^{-4} ppm) and nitrogen contents (47 ppm) and isotopic ratio of nitrogen in 14313.




Figure 8: Ratio of clast abundance in Apollo 14 breccia samples with 14313 (Simonds et al. 1977).

Mineralogical Mode for 14313

Drozd et al. 1977	
Mineral fragments	30.9 %
Lithic fragments	19.6
Colored glass	11
Agglutinate glass	36.8
Colorless glass	0.6
Chondrules	0
Devitrified glass	1

Mineralogical Mode for 14313

	Simonds et al. 1977	Drozd et al. 1976	Simon et al. 1989
Matrix Clasts	72 %		55.1
Plagioclase	4		5.8
Mafic	1		6.7
Breccia	11.5		~20
Glass	2	12.6	8.1
Granulite	9		2.6
Basalt	1		0.5
Agglutinate		36.8	7.6




Figure 9: Basaltic or "micronorite" clast in 14313. Scale bar is 100 microns. Floran et al. 1972.




Figure 11: Ar-Ar release pattern with no age information Alexander and Kahl (1974).




Figure 10: Carbon content of Apollo 14 breccias with values for 14313 (Moore et al. 1972).





Figure 12: Measured seismic wave velocity as function of pressure for breccia 14313 (Mizutani et al. 1972).

Table 1. Chemical composition of 14313 .


reference weight	Boynton 75 262 259 mg		Helmke72		Simon 89 Hughes 73 156 mg			
SiO2 %								
TiO2					1.68	(a)		
Al2O3	17.38		(a)		18.4	(a)		
FeO	8.6	10.7	(a)		10.3	(a)		
MnO	0.13		(a)	0.13	(a)	0.123	(a)	
MgO					9.4	(a)		
CaO	9.8	10.6	(a)		11	(a)		
Na2O	0.77		(a)		0.77	(a)		
K2O					0.53	(a)		
P2O5								
S %								
sum								
Sc ppm	19.2	22	(a)	24.6	(a)	21.4	(a)	
V					51	(a)		
Cr	1200	1370	(a)	2450	(a)	1290	(a)	
Co	30	36	(a)	31.5	(a)	32	(a)	
Ni	338		(b)	273	(a)	390	(a)	
Cu								
Zn	35		(b)	56	(a)			
Ga	8.09		(b)	11	(a)			
Ge ppb	700		(b)					
As								
Se								
Rb					13	(a)		
Sr					110	(a)		
Y								
Zr					750	(a)		
Nb								
Mo								
Ru								
Rh								
Pd ppb								
Ag ppb							16	12 (b)
Cd ppb	280		(b)					
In ppb	24		(b)					
Sn ppb								
Sb ppb								
Te ppb								
Cs ppm					0.75	(a)		
Ba	690	760	(a)		810	(a)		
La	64	70	(a)	65.4	(a)	67	(a)	
Ce	160	190	(a)	171	(a)	176	(a)	
Pr								
Nd				114	(a)	108	(a)	
Sm	31	31	(a)	30.8	(a)	28.7	(a)	
Eu	2.46	2.81	(a)	2.65	(a)	2.8	(a)	
Gd				40	(a)			
Tb	5	6.2	(a)	6.3	(a)	6	(a)	
Dy	33		(a)	38	(a)	36	(a)	
Ho				8	(a)			
Er				22	(a)			
Tm								
Yb	20	24	(a)	21.8	(a)	20.1	(a)	
Lu	2.6	3.2	(a)	3.17	(a)	2.65	(a)	
Hf	19	20	(a)	21	(a)	21.3	(a)	
Ta	3		(a)		2.9	(a)		
W ppb								
Re ppb							1.21	0.51 (b)
Os ppb							16.8	7.1 (b)
Ir ppb	10.3		(b)		7	(a)	18	8.1 (b)
Pt ppb								
Au ppb	5.1		(b)		7.2	(a)	8	4.8 (b)
Th ppm	11	13	(a)		11.9	(a)		
U ppm					3.1	(a)		



technique: (a) INAA, (b) RNAA



C Meyer
2007

14313
144 g





References for 14313

- Alexander E.C. and Kahl S.B. (1974) ^{40}Ar - ^{39}Ar studies of lunar breccias. Proc. 5th Lunar Sci. Conf. 1353-1373.
- Basford J.R., Dragon J.C., Pepin R.O., Coscio M.R. and Murthy V.R. (1972) Krypton and Xenon in lunar fines. Proc. 4th Lunar Sci. Conf. 1915-1955.
- Becker R.H. and Clayton R.N. (1975) Nitrogen abundances and isotopic compositions in lunar samples. Proc. 6th Lunar Sci. Conf. 2131-2149.
- Behrmann C.J., Drozd R.J. and Hohenberg C.M. (1973) Extinct lunar radio-activities: Xenon from ^{244}Pu and ^{129}I in Apollo 14 breccias. Earth Planet. Sci. Lett. 17, 446-455.
- Boynton W.V., Baedeker P.A., Chou C-L., Robinson K.L. and Wasson J.T. (1975) Mixing and transport of lunar surface materials: Evidence obtained by the determination of lithophile, siderophile and volatile elements. Proc. 6th Lunar Sci. Conf. 2241-2259.
- Carlson I.C. and Walton W.J.A. (1978) **Apollo 14 Rock Samples**. Curators Office. JSC 14240
- Chao E.C.T., Minkin J.A. and Best J.B. (1972) Apollo 14 breccias: General characteristics and classification. Proc. 3rd Lunar Sci. Conf. 645-659.
- Delano J.W. (1987) Apollo 14 regolith breccias: Different glass populations and their potential for charting space/time variations. Proc. 18th Lunar Planet Sci. Conf. 59-65. Lunar Planetary Institute, Houston.
- Drozd R.J., Kennedy B.M., Morgan C.J., Podosek F.A. and Taylor G.J. (1976) The excess fission xenon problem in lunar samples. Proc. 7th Lunar Sci. Conf. 599-623.
- Dunlap D.J., Gose W.A., Pearce G.W. and Strangway D.W. (1973) Magnetic properties and granulometry of metallic iron in lunar breccia 14313. Proc. 4th Lunar Sci. Conf. 2977-2990.
- Floran R.J., Cameron K.L., Bence A.E. and Papike J.J. (1972) Apollo 14 breccia 14313: a mineralogic and petrologic report. Proc. 3rd Lunar Sci. Conf. 661-671.
- Fruland R.M. (1983) Regolith Breccia Workbook. JSC 19045
- Gose W.A., Pearce G.W., Strangway D.W. and Larson E.E. (1972) Magnetic properties of Apollo 14 breccias and their correlation with metamorphism. Proc. 3rd Lunar Sci. Conf. 2387-2395.
- Graf H., Shirck J., Sun S and Walker R. (1973) Fission track astrology of three Apollo 14 gas-rich breccias. Proc. 4th Lunar Sci. Conf. 2145-2155.
- Helmke P.A., Haskin L.A., Korotev R.L. and Ziege K.E. (1972) Rare earths and other trace elements in Apollo 14 samples. Proc. 3rd Lunar Sci. Conf. 1275-1292.
- Hughes T.C., Keays R.R. and Lovering J.F. (1973) Siderophile and volatile trace elements in Apollo 14, 15 and 16 rocks and fines: Evidence for extralunar component and Ti-, Au- and Ag-enriched rocks in the ancient lunar crust (abs). Lunar Sci. IV, 400-402. Lunar Planetary Institute, Houston.
- Juan V.C., Chen J.C., Huang C.K., Chen P.Y. and Wang Lee C.M. (1972) Petrology and chemistry of some Apollo 14 lunar samples. Proc. 3rd Lunar Sci. Conf. 687-705.
- King E.A., Butler J.C. and Carman M.F. (1972) Chondrules in Apollo 14 samples and size analyses of Apollo 14 and 15 fines. Proc. 3rd Lunar Sci. Conf. 673-686.
- LSPET (1971) Preliminary examination of lunar samples from Apollo 14. Science 173, 681-693.
- Mizutani H., Fujii N., Hamano Y. and Osako M. (1972) Elastic wave velocities and thermal diffusivities of Apollo 14 rocks. Proc. 3rd Lunar Sci. Conf. 2557-2564.
- Moore C.B., Lewis C.F., Cripe J., Delles F.M., Kelly W.R. and Gibson E.K. (1972) Total carbon, nitrogen and sulfur in Apollo 14 lunar samples. Proc. 3rd Lunar Sci. Conf. 2051-2058.
- Morrison D.A., McKay D.S., Heiken G.H. and Moore H.J. (1972) Microcraters on lunar rocks. Proc. 3rd Lunar Sci. Conf. 2767-2791.
- Pearce G.W., Strangway D.W. and Gose W.A. (1972) Remanent magnetism of the lunar surface. Proc. 3rd Lunar Sci. Conf. 2449-2464.
- Ruzicka A., Snyder G.A. and Taylor L.A. (2000) Crystal-bearing lunar spherules: Impact melting of the Moon's crust and implications for the origin of meteoritic chondrules. Meteoritics & Planet. Sci. 35, 173-192.
- Simon S.B., Papike J.J., Shearer C.K., Hughes S.S. and Schmitt R.A. (1989) Petrology of Apollo 14 regolith breccias and ion microprobe studies of glass beads. Proc. 19th Lunar Planet. Sci. Conf. 1-17. Lunar Planetary Institute, Houston.
- Simonds C.H., Phinney W.C., Warner J.L., McGee P.E., Geeslin J., Brown R.W. and Rhodes J.M. (1977) Apollo 14 revisited, or breccias aren't so bad after all. Proc. 8th Lunar Sci. Conf. 1869-1893.

Sutton R.L., Batson R.M., Larson K.B., Schafer J.P., Eggleton R.E. and Swann G.A. (1971) Documentation of the Apollo 14 samples. U.S. Geological Survey, Rpt. 32.

Sutton R.L., Hait M.H. and Swann G.A. (1972) Geology of the Apollo 14 landing site. Proc. 3rd Lunar Sci. Conf. 27-38.

Swann G.A., Trask N.J., Hait M.H. and Sutton R.L. (1971a) Geologic setting of the Apollo 14 samples. Science 173, 716-719.

Swann G.A., Bailey N.G., Batson R.M., Eggleton R.E., Hait M.H., Holt H.E., Larson K.B., Reed V.S., Schaber G.G., Sutton R.L., Trask N.J., Ulrich G.E. and Wilshire H.G. (1977) Geology of the Apollo 14 landing site in the Fra Mauro Highlands. U.S.G.S Prof. Paper 880.

Swann G.A., Bailey N.G., Batson R.M., Eggleton R.E., Hait M.H., Holt H.E., Larson K.B., McEwen M.C., Mitchell E.D., Schaber G.G., Schafer J.P., Shepard A.B., Sutton R.L., Trask N.J., Ulrich G.E., Wilshire H.G. and Wolfe E.W. (1972) 3. Preliminary Geologic Investigation of the Apollo 14 landing site. In Apollo 14 Preliminary Science Rpt. NASA SP-272. pages 39-85.

Warner J.L. (1972) Metamorphism of Apollo 14 breccias. Proc. 3rd Lunar Sci. Conf. 623-643.

Warren P.H. (1993) A concise compilation of petrologic information on possibly pristine nonmare Moon rocks. Am. Mineral. 78, 360-376.

Warren P.H., Taylor G.J., Keil K., Kallemeyn G.W., Shirley D.N. and Wasson J.T. (1983) Seventh foray: Whitlockite-rich lithologies, a diopside-bearing troctolitic anorthosite, ferroan anorthosites and KREEP. Proc. 14th Lunar Planet. Sci. Conf., J. Geophys. Res. B151-B164.

Wentworth S.J. and McKay D.S. (1991) Apollo 14 glasses and the origin of lunar soils. Proc. 21st Lunar Planet. Sci. Conf. 185-192. Lunar Planetary Institute, Houston.

Williams R.J. (1972) The lithification of metamorphism of lunar breccias. Earth Planet. Sci. Lett. 16, 250-256.

Wilshire H.G. and Jackson E.D. (1972) Petrology and stratigraphy of the Fra Mauro Formation at the Apollo 14 site. U.S. Geol. Survey Prof. Paper 785.