

65785

revised

Impact-melt with Pink Spinel Troctolite clast

5.16 grams




Figure 1: Photo of rake sample 65785 showing clast of pink spinel troctolite (PST) imbedded in recrystallized impact-melt. Scale in mm. NASA S72-48821.

Introduction

This small rake sample contains a coarse-grained pink spinel troctolite clast in a crystalline feldspathic impact melt rock (figure 1). The impact melt has been dated at 3.97 b.y. by one group and 3.83 b.y. by another.

Petrography

Dowty et al. (1974b) describe the pink spinel troctolite clast (figure 2). It is modally 65% plagioclase (An_{98}), 30% olivine (Fo_{85}) and 5% Mg-Al spinel. The pink spinel occurs as one large crystal (1 mm). Irregular plagioclase grains (0.1 – 1 mm) are surrounded by one large olivine grain. Minor phases are pyroxene,

ilmenite, armalcolite, rutile, metal, troilite, whitlockite, farringtonite and K-feldspar (Warner et al. 1976).

Warner et al. (1976) describe the crystalline impact-melt rock that surrounds the clast. In it the plagioclase (An_{95}) occurs predominantly as laths about 0.2 mm long optically within olivine (Fo_{79}) and minor pyroxene grains. The fine-grained portion also has trace spinel, chromite, armalcolite, whitlockite, apatite, zirkelite and schreibersite as minor phases.

Keil et al. (1975) suggested that the melt-rock portion of 65785 may be a mixture of the troctolite lithology




Figure 2: Thin section photomicrograph of pink spinel troctolite clast in 65785 (Dowty et al. 1974). Large spinel grain at bottom. Width of field about 2 mm.

with KREEP. Ryder and Norman (1980) summarize the results. Compositional variation of Apollo 16 impact-melt rocks is discussed by Korotev (1994).

Mineralogy

Olivine: The olivine occurs as a large grain surrounding and including the other phases in the thin section of the pink spinel troctolite clast. It is slightly more mafic (Fo_{85}) than the olivine found in the surrounding matrix (Fo_{80}).

Farringtonite: Dowty et al. (1974b) and Warner et al. (1976) reported a rare phosphate ($Mg_3(PO_4)_2$) in the pink spinel troctolite clast.

Pyroxene: The pyroxene in 65785 is mafic in composition (figures 4 and 5).

Plagioclase: Plagioclase is An95-98.

Mg-Al Spinel: The large pink spinel grain in 65785 is zoned in Cr content (2.6 to 12.6%), increasing outward from the center (Dowty et al. 1974b).

Armalcolite: Dowty et al. (1974) determined the composition of armalcolite grains and reported on unpublished Pb-Pb age dating by ion microprobe analysis.

Metallic iron: Dowty et al. (1974b) found Ni = 2-25%, Co = 0.5-1.5%




Figure 3: Plagioclase and pyroxene composition diagram for 65785.




Figure 4: Pyroxene and olivine composition of troctolite clast (PST) in 65785 (from Dowty et al. 1974).




Figure 5: Pyroxene and olivine composition of melt-rock portion of 65785 (Warner et al. 1976).




Figure 6: Composition of matrix portion of 65785 (Murali et al. 1977).

Chemistry

The pink spinel troctolite composition was determined by broad beam electron probe analysis (Warner et al. 1976). Murali et al. (1977) reported the trace element analysis of the surrounding impact-melt (table 1, figure 6).

Radiogenic age dating

Schaeffer and Schaeffer (1977) determined the Ar/Ar plateau age of the feldspathic matrix of 65785 (figure 7). Dowty et al. (1974) reported ion probe Pb-Pb age data by Anderson and Hinthorne for three armalcolite grains in the troctolite clast.

Norman et al. (2006) reported an Ar/Ar plateau age of 3.83 ± 0.02 b.y. (figure 8).

Cosmogenic isotopes and exposure ages

Eldridge et al. (1974) found the cosmic ray induced activity for $^{22}\text{Na} = 45$ dpm/kg. and $^{26}\text{Al} = 59$ dpm/kg.




Figure 7: Argon release pattern for 65785 (Schaeffer and Schaeffer 1977).




Figure 8: Argon release pattern for 65785 (Norman et al. 2006).

Summary of Age Data for 65785


	Pb/Pb	Ar/Ar
Schaeffer and Schaeffer 1977		3.97 ± 0.02 b.y.
Anderson and Hinthorne	4.0	in Dowty et al. 1974
Norman et al. 2006		3.83 ± 0.02

Note: old decay constant for K.


Schaeffer and Schaeffer (1977) determined a cosmic ray exposure age of 271 m.y.

Processing

There are 4 thin sections of 65785.



*Figure 9: Thin section
65785,22. 2 mm
across*



References for 65785

- Butler P. (1972a) Lunar Sample Information Catalog Apollo 16. Lunar Receiving Laboratory. MSC 03210 Curator's Catalog. pp. 370.
- Dowty E., Keil K. and Prinz M. (1974a) Igneous rocks from Apollo 16 rake samples. *Proc. 5th Lunar Sci. Conf.* 431-445.
- Dowty E., Green J.A., Hlava P.F., Keil K., Moore R.B., Nehru C.E., Prinz M. and Warner R.D. (1976) Electron microprobe analyses of minerals from Apollo 16 rake samples. *Inst. Meteoritics Spec. Publ. No 14*, 141 pp.. Univ. New Mex. ABQ
- Keil K., Dowty E., Prinz M. and Bunch T.E. (1972) Description, classification and inventory of 151 Apollo 16 rake samples from the LM area and station 5. Curator's Catalog, JSC.
- Keil K., Warner R.D., Prinz M. and Dowty E. (1975) Rocks 60618 and 65785: Evidence for admixture of KREEP in lunar impact melts. *Geophys. Res. Lett.* **2**, 369.
- Korotev R.L. (1994) Compositional variation in Apollo 16 impact melt breccias and inferences for the geology and bombardment history of the central highlands of the Moon. *Geochim. Cosmochim. Acta* **58**, 3931-3969.
- LSPET (1973b) The Apollo 16 lunar samples: Petrographic and chemical description. *Science* **179**, 23-34.
- LSPET (1972c) Preliminary examination of lunar samples. *In Apollo 16 Preliminary Science Report. NASA SP-315*, 7-1—7-58.
- McKinley J.P., Taylor G.J., Keil K., Ma M.-S. and Schmitt R.A. (1984) Apollo 16: Impact sheets, contrasting nature of the Cayley Plains and Descartes Mountains, and geologic history. *Proc. 14th Lunar Planet. Sci. Conf.* in *J. Geophys. Res.* **89**, B513-B524.
- Norman M.D., Duncan R.A. and Huard J.J. (2006) Identifying impact events within the lunar cataclysm from ⁴⁰Ar-³⁹Ar ages and compositions of Apollo 16 impact melt rocks. *Geochim. Cosmochim. Acta* **70**, 6032-6049.
- Reimold W.U., Nyquist L.E., Bansal B.M., Wooden J.L., Shih C.-Y., Wiesmann H. and Mackinnon I.D.R. (1985) Isotope analysis of crystalline impact-melt rocks from Apollo 16 stations 11 and 13. North Ray Crater. *Proc. 15th Lunar Planet. Sci. Conf.* in *J. Geophys. Res.* **90**, C431-C448.
- Ryder G. and Norman M.D. (1980) Catalog of Apollo 16 rocks (3 vol.). Curator's Office pub. #52, JSC #16904
- Sutton R.L. (1981) Documentation of Apollo 16 samples. *In Geology of the Apollo 16 area, central lunar highlands.* (Ulrich et al.) U.S.G.S. Prof. Paper 1048.
- Warner R.D., Dowty E., Prinz M., Conrad G.H., Nehru C.E. and Keil K. (1976c) Catalog of Apollo 16 rake samples from the LM area and station 5. *Spec. Publ. #13*, UNM Institute of Meteoritics, Albuquerque. 87 pp.
- Warren P.H. (1993) A concise compilation of petrologic information on possibly pristine nonmare Moon rocks. *Am. Mineral.* **78**, 360-376.
- Wiesmann H. and Hubbard N.J. (1975) A compilation of the Lunar Sample Data Generated by the Gast, Nyquist and Hubbard Lunar Sample PI-Ships. Unpublished. JSC