

70017
Ilmenite Basalt
2957 grams



Figure 1: Photo of lunar basalt 70017 showing vugs and vesicles. Sample is about 10 cm across. Cube is 1 cm. NASA# S73-15723.

Introduction

Lunar sample 70017 is a vesicular, medium-grained, high-Ti basalt that was collected from the area near the Lunar Lander (figure 1). It was one of the last rocks collected from the moon and the astronauts made a speech about it referring to “the children of the world”. Subsequently, small pieces of it were distributed widely. It has also been the subject of many science investigations (briefly summarized below).

70017 has a crystallization age about 3.7 b.y. and an exposure age about 220 m.y.

Petrography

Longhi et al. (1974) describe 70017 as “a medium-grained hypidiomorphic granular high-Ti basalt with textural relations suggesting relatively slow cooling. Large, equant, subhedral grains of clinopyroxene (1-2




Figure 2: Thin section 70017,111 showing “chains” of ilmenite. Field of view 1.5 cm.




Figure 3. Three views of same area of thin section 70017,125 (transmitted light, crossed polarizers, and reflected light). Field of view is 2.5 mm. NASA # S79-26723-5.

mm) enclose embayed ilmenite (figure 3). Plagioclase (up to 2 mm) is anhedral and poikilitically encloses clinopyroxene, olivine and ilmenite. Cristobalite is present in interstices between clinopyroxene, plagioclase and ilmenite.”



Figure 4: Partially polarized light photo of polysynthetic twinning in plagioclase poikilitically enclosing equant olivine and pyroxene crystals in 70017.

Roedder and Weiblen (1975) studied melt inclusions in ilmenite in 70017 and other Apollo 17 rocks and found evidence for silicate liquid immiscibility.

Nord et al. (1974) determined that 70017 had a single-stage cooling history (<2 deg./hr.), by carefully studying the defect structure of pyroxene and plagioclase. Usselman et al. (1975) also calculated the cooling rate.

Mineralogy

Olivine: Olivine (Fo₆₆₋₆₉) is found in the cores of the large clinopyroxene grains and in plagioclase (figure 4).

Pyroxene: Pyroxene is abundant in 70017. Large pyroxene grains are zoned from subcalcic augite to pigeonite to Fe-rich (figure 6). Pyroxene cores contain up to 3.5% TiO₂ and 4.5% Al₂O₃ (Brown et al. 1973, Hodges and Kushiro 1974 and Longhi et al. 1974). Sung et al. (1974) studied the ratio of ³⁺Ti/⁴⁺Ti in pyroxene – perhaps 30% is ³⁺Ti!



Figure 5: Reflected light photo of polished thin section illustrating habit of ilmenite in 70017.

Plagioclase: Longhi et al. (1974) measured plagioclase as An₆₉₋₈₈.

Opaques: Ilmenite is abundant and found occurring in long “chains” (figure 2). It has exsolution of chromite and rutile (figure 5). Experiments show that armalcolite was an early phase, but it has mostly converted to ilmenite. Some relict armalcolite is found in pyroxene cores (figure 7). Small grains of chromite are reported in olivine (Hodges and Kushiro 1974). El Goresy and Ramdohr (1975) studied the opaque minerals in 70017 in order to determine the nature of subsolidus reduction




Figure 6: Pyroxene and olivine composition for mare base 70017 (data copied from Longhi et al. 1974 and Hodges and Kushiro 1974).

producing elemental iron. Analyses of armalcolite in 70017 are given in table 2.

Silica: Silica found in 70017 is cristobalite (Hodges and Kushiro 1974, Roedder and Weiblen 1975) and has a characteristic cracked texture (figure 8).

Mesostasis: A very complex mesostasis of glass, silica, troilite-iron, whitlockite and trace tranquillityite occurs in this rock.

Chemistry

The chemical composition of 70017 is given in table 1 and figures 9 and 10. It is typical of the other basalts returned from Taurus-Littrow (figure 11) and is classified as a group B basalt (Paces et al. 1991). The chemical composition of high and low K melt inclusions is found in Roedder and Weiblen (1975).

Radiogenic age dating

Nyquist et al. (1975) and Schaeffer et al. (1977) have dated 70017 at 3.67 b.y. Note that the “ilmenite separates” are found to contain the most Rb (figure 12). The “ages” determined by Phinney et al. (1975) and Mattinson et al. (1977) were not convincing. Tilton

Mineralogical Mode for 70017

	Longhi et al. 1974	Brown et al. 1975	Roedder, Weiblen 1975
Olivine	1 %	0.9	0.4
Pyroxene	50	49.3	57.6
Plagioclase	26	25.4	19.8
Opaques	22	22.8	19.2
Fe	tr.		tr.
FeS	tr.		
Mesostasis	tr.	0.3	1.4
Silica		1.3	1.6




Figure 7: Reflected light photo of tan, barrel-shaped armalcolite in 70017. It is hexagonal in cross section (center of photo), but tapered in the elongate direction.




Figure 8: Reflected light photo of silica in 70017.

and Chen (1979) also reported U/Pb data for 70017. Additional isotopic data is found in Unruh et al. (1984).

Cosmogenic isotopes and exposure ages

Phinney et al. (1975) determined an exposure age of 220 ± 20 m.y. for 70017.

Yokoyama et al. (1974) report that 70017 was apparently saturated with ^{26}Al .

Other Studies

Rutherford et al. (1974) studied the “liquid line of descent” for the cooling of a basalt with the composition of 70017 and found experimental evidence for “liquid immiscibility”. Sato (1976) determined the *in situ* oxygen fugacity as a function of temperature (figure 13).

Petrowski et al. (1974) determined the concentration and isotopic composition of sulfur and carbon. Mayeda et al. (1974) reported the isotopic composition of oxygen.

Huffman et al. (1974) measured the Mössbauer spectra of 70017 (figure 14) and determined the amount of metallic iron (0.1%). Brecher et al. (1974), Nagata (1974), Stephenson et al. (1974) and Schwerer and Nagata (1976) studied the magnetic properties of 70017.

Hapke et al. (1978) and Osborne et al. (1978) used powdered rock samples of 70017 to calibrate spectra that might be obtained from orbit.

Table 2: Analysis of armalcolite in 70017.

from El Goresy et al. 1974

SiO ₂	0.33	0.3	0.32	0.3
Al ₂ O ₃	1.72	1.77	1.6	1.57
TiO ₂	70.44	71.61	71.69	71.7
Cr ₂ O ₃	1.63	1.69	1.46	1.39
FeO	18.43	16.3	15.24	15.5
MnO	0.08	0.09	0.1	0.1
MgO	5.47	6.63	7.16	7.41
CaO	0.57	0.44	0.4	0.45
V ₂ O ₃	0.18	0.27	0.39	0.38
total	98.85	99.1	98.36	98.8




Figure 9: 70017 is a typical Apollo 17 basalt.




Figure 10: Normalized rare-earth-element pattern of 70017 (isotope dilution data from two different labs).

Processing

Figure 15 shows the initial cutting of 70017.

Neal and Taylor (1993) provide an extensive review of the work done on 70017 in their catalog. Sixty-four thin sections were prepared from 70017. Thin sections of this rock are featured in the Lunar Petrographic Educational Thin Section Set (Meyer 2003).

Small pieces of this rock were given to many countries of the world (70017 has been called the “children of the world rock”). A piece of this rock is on public display at the Virginia Air and Space Center, Hampton Virginia (figure 16).

List of Photo #s for 70017

- S73-15719 – 724 color
- S73-21892 – 907 B&W
- S73-28689 processing
- S76-21593
- S79-26723 – 725 TS
- S80-40284 – 287 TS
- S80-42343 TS
- S92-32791 – 793 display
- S92-33835 – 837




Figure 11: Trace element diagram used to distinguish Apollo 17 basalt flows.




Figure 12: Rb-Sr isochron for 70017 (from Nyquist et al. 1974).

Summary of Age Data for 70017

	Ar-Ar	Rb-Sr	U-Pb
Phinney et al. 1975	3.80 ± 0.03 b.y.		
Nyquist et al. 1975		3.67 ± 0.18	
Mattinson et al. 1977			3.7 b.y.
Schaeffer et al. 1977	3.67 ± 0.12		

Caution: Old decay constants




Figure 13: Oxygen fugacity vs. temperature for 70017 (from Sato 1976).




Figure 14: Moessbauer spectra of 70017 (from Huffman et al. 1974)



Figure 16: Display sample 70017,138.

Table 1. Chemical composition of 70017.

reference weight	Rose 74		Brunfelt 74	Miller 74	Nava 74	Rhodes 74	Duncan74	Philpotts74	Shih 75	Dickinson 89
				Hughes85		Unruh84			Wiesmann75	
SiO2 %	38.8	38.68	(a)	38.52	38.8	38.07	(f) 38.37	(f)		
TiO2	12.84	13.75	(a) 13.58	12.21	12.44	13.1	(f) 12.83	(f)		
Al2O3	8.54	7.4	(a) 9.47	9.07	9.73	8.79	(f) 8.78	(f)		
FeO	18.12	18.77	(a) 18.32	18.19	17.6	18.07	(f) 18.71	(f)		17.5 (b)
MnO	0.24	0.25	(a) 0.254	0.233	0.232	0.27	(f) 0.247	(f)		
MgO	10.16	10.45	(a) 9.13	11.95	9.89	9.81	(f) 9.41	(f)		
CaO	10.56	10.05	(a) 11.48	10.36	10.04	10.3	(f) 10.43	(f)		7.14 13.4 (b)
Na2O	0.33	0.34	(a) 0.405	0.405	0.43	0.4	(f) 0.43	(f)	0.31	0.32 0.43 (b)
K2O	0.07	0.07	(a) 0.044		0.036	0.04	(f) 0.047	(f)	0.038	(e)
P2O5	0.04	0.04	(a)		0.048	0.05	(f) 0.052	(f)		
S %						0.15	(f) 0.175	(f)		
sum						99.05				
Sc ppm	80	77	(a) 87	(b)					82.7	(e) 75 78 (b)
V	98	80	(a) 156	(b)			146	(f)		288 (b)
Cr	3350	3350	(a) 3550	(b)	3080		3948	(f)	3490	(e) (b)
Co	32	32	(a) 20.6	(b)			18	(f)	14.5	(e) 132 22 (b)
Ni	<1	24	(a) <10	(b)			<3	(f)		
Cu	28	84	(a) 2.8	(b)			<3	(f)		
Zn	<4	<4	(a) 2	(b)			<2	(f)		
Ga	5.8	5.4	(a) 3.1	(b)						21 (b)
Ge ppb									1.7	1.9 (b)
As										
Se										
Rb	0.9	0.7	(a) 0.4	(b)			1.2	(f) 0.28	0.299	(e) (b)
Sr	217	155	(a) 127	(b)			166	(f) 168	153	(e) 172 306 (b)
Y	94	100	(a)				71.2	(f) 32		
Zr	254	250	(a)	304	(b)		218	(f) 223	177	(e) 138 (b)
Nb	23	18	(a)				18.5	(f)		
Mo										
Ru										
Rh										
Pd ppb										
Ag ppb										
Cd ppb										
In ppb										
Sn ppb										
Sb ppb										
Te ppb										
Cs ppm			0.3	(b)						
Ba	250	180	(a) 55	(b)			83	(f) 43	45.8	(e) 78 68 (b)
La	<10	<10	(a) 4.11	(b)					3.99	(e) 4.6 4.4 (b)
Ce			13.5	(b)				10.7	11.3	(e) 16 15 (b)
Pr										
Nd						17.5	(e)	12.1	13.2	(e) 20 14 (b)
Sm			7.53	(b)		7.4	(e)	5.13	5.67	(e) 6.7 71 (b)
Eu			1.77	(b)				1.62	1.49	(e) 1.6 1.7 (b)
Gd									9.05	(e)
Tb			1.77	(b)						2.5 2 (b)
Dy			13.8	(b)				10.2	10.7	(e)
Ho										
Er								6.31	6.46	(e) 0.71 (b)
Tm										
Yb	7.7	8.3	(a) 6.3	(b)				6.25	5.98	(e) 8.2 6.9 (b)
Lu			1.15	(b)		1	(e)	0.954	0.933	(e) 1.3 1.1 (b)
Hf			8	(b) 7.8	(b)	7	(e)			7.4 6 (b)
Ta			1.55	(b)						1.8 1.5 (b)
W ppb			0.075	(b)						
Re ppb										
Os ppb										
Ir ppb										
Pt ppb										
Au ppb										
Th ppm			0.17	(b)					0.198	(e) 4.8 0.14 (b)
U ppm			0.088	(b)					0.06	(e)

technique: (a) combined XRF, OES, chemical, (b) INAA, (e) IDMS, (f) XRF





Figure 15: Sample 70017 after initial sawing. Note high proportion of vesicles. Small cube is 1 cm.
NASA# S73-28689



References for 70017

- Boyett M. and Carlson R.W. (2007) A highly depleted moon or a non-magma ocean origin for the lunar crust? *Earth Planet. Sci. Lett.* 262, 505-516.
- Brecher A. (1974) Inferences from comparative magnetic studies of some Apollo 17 basalts, breccias and soils (abs). *Lunar Sci.* V, 83-85. Lunar Planetary Institute, Houston.
- Brown G.M., Peckett A., Emeleus C.H., Phillips R. and Pinsent R.H. (1975a) Petrology and mineralogy of Apollo 17 mare basalts. *Proc. 6th Lunar Sci. Conf.* 1-13.
- Brunfelt A.O., Heier K.S., Nilssen B., Steinnes E. and Sundvoll B. (1974) Elemental composition of Apollo 17 fines and rocks. *Proc. 5th Lunar Sci. Conf.* 981-990.
- Butler P. (1973) **Lunar Sample Information Catalog Apollo 17.** Lunar Receiving Laboratory. MSC 03211 Curator's Catalog. pp. 447.
- Dickinson T., Taylor G.J., Keil K. and Bild R.W. (1989) Germanium abundances in lunar basalts: Evidence of mantle metasomatism. *Proc. 19th Lunar Planet. Sci.* 189-198. Lunar Planetary Institute, Houston
- Duncan A.R., Erlank A.J., Willis J.P., Sher M.K. and Ahrens L.H. (1974a) Trace element evidence for a two-stage origin of some titaniferous mare basalts. *Proc. 5th Lunar Sci. Conf.* 1147-1157.
- Hodges F.N. and Kushiro I. (1974a) Apollo 17 petrology and experimental determination of differentiation sequences in model Moon compositions. *Proc. 5th Lunar Sci. Conf.* 505-520.
- El Goresy A., Ramdohr P., Medenbach O. and Bernhardt H.-J. (1974a) Taurus-Littrow TiO₂-rich basalts: Opaque mineralogy and geochemistry. *Proc. 5th Lunar Sci. Conf.* 627-652.
- El Goresy A., Ramdohr P., Medenbach O. and Bernhardt H.-J. (1974b) Taurus-Littrow crystalline rocks: Opaque mineralogy and geochemistry (abs). *Lunar Sci.* V, 209-211. Lunar Planetary Institute, Houston
- El Goresy A. and Ramdohr P. (1975a) Subsolvus reduction of lunar opaque oxides: Textures, assemblages, geochemistry, and evidence for a late-stage endogenic gaseous mixture. *Proc. 6th Lunar Sci. Conf.* 729-745.
- El Goresy A. and Ramdohr P. (1975b) Subsolvus reduction of lunar opaque oxides: Evidence, assemblages, geochemical relevance, and evidence for a late-stage reducing gaseous mixture (abs). *Lunar Sci.* VI, 245-247. Lunar Planetary Institute, Houston
- El Goresy A. and Ramdohr P. (1975c) Taurus-Littrow TiO₂-rich basalts: Opaque mineralogy and geochemistry (abs). *Lunar Sci.* VI, 248-250. Lunar Planetary Institute, Houston
- Green D.H., Ringwood A.E., Hibberson W.O. and Ware N.G. (1975a) Experimental petrology of Apollo 17 mare basalts. *Proc. 6th Lunar Sci. Conf.* 871-893.
- Hapke B.W., Partlow W.D., Wagner J.K. and Cohen A.J. (1978) Reflectance measurements of lunar materials in the vacuum ultraviolet. *Proc. 9th Lunar Planet. Sci. Conf.* 2935-2947.
- Huffman G.P., Schwerer F.C., Fisher R.M. and Nagata T. (1974a) Iron distributions and metallic-ferrous ratios for Apollo lunar samples: Mossbauer and magnetic analyses. *Proc. 5th Lunar Sci. Conf.* 2779-2794.
- Hughes S.S. and Schmitt R.A. (1985) Zr-Hf-Ta fractionation during lunar evolution. *Proc. 16th Lunar Planet. Sci. Conf.* D31 in *J. Geophys. Res.* 90.
- Lindstrom M.M. and Haskin L.A. (1978) Causes of compositional variations within mare basalt suites. *Proc. 9th Lunar Planet. Sci. Conf.* 465-486.
- Longhi J., Walker D., Grove T.L., Stolper E.M. and Hays J.F. (1974) The petrology of the Apollo 17 mare basalts. *Proc. 5th Lunar Sci. Conf.* 447-469.
- LSPET (1973) Apollo 17 lunar samples: Chemical and petrographic description. *Science* 182, 659-672.
- LSPET (1973) Preliminary Examination of lunar samples. Apollo 17 Preliminary Science Rpt. NASA SP-330. 7-1 – 7-46.
- Mattinson J.M., Tilton G.R., Todt W. and Chen J.H. (1977) Lead isotope studies of mare basalt 70017. *Proc. 8th Lunar Sci. Conf.* 1473-1487.
- Mayeda T.K., Shearer J. and Clayton R.N. (1975) Oxygen isotope fractionation of Apollo 17 rocks. *Proc. 6th Lunar Sci. Conf.* 1799-1802.
- McGee P.E., Simonds C.H., Warner J.L. and Phinney W.C. (1979) Introduction to the Apollo Collections:
- Meyer C. (1994) Catalog of Apollo 17 rocks. Vol. 4 North Massif
- Meyer C. (1987, 2003) The Lunar Petrographic Thin Section Set. Curatorial Branch Publication No. 76. JSC. Houston
- Miller M.D., Pacer R.A., Ma M.-S., Hawke B.R., Lookhart G.L. and Ehmann W.D. (1974) Compositional studies of

- the lunar regolith at the Apollo 17 site. Proc. 5th Lunar Sci. Conf. 1079-1086.
- Muehlberger et al. (1973) Documentation and environment of the Apollo 17 samples: A preliminary report. *Astrogeology* 71 322 pp superceded by *Astrogeology* 73 (1975) and by Wolfe et al. (1981)
- Muehlberger W.R. and many others (1973) Preliminary Geological Investigation of the Apollo 17 Landing Site. *In Apollo 17 Preliminary Science Report*. NASA SP-330.
- Nagata T., Sugiura N., Fisher R.M., Schwerer F.C., Fuller M.D. and Dunn J.R. (1974a) Magnetic properties of Apollo 11-17 lunar materials with special reference to effects of meteorite impact. Proc. 5th Lunar Sci. Conf. 2827-2839.
- Nagata T., Sugiura N., Fisher R.M., Schwerer F.C., Fuller M.D. and Dunn J.R. (1974b) Magnetic properties and natural remanent magnetization of Apollo 16 and 17 lunar samples (abs). *Lunar Sci.* V, 540-542. Lunar Planetary Institute, Houston
- Nagata T., Fisher R.M., Schwerer F.C., Fuller M.D. and Dunn J.R. (1975a) Effects of meteorite impact on magnetic properties of Apollo lunar materials. Proc. 6th Lunar Sci. Conf. 3111-3122.
- Nava D.F. (1974a) Chemical compositions of some soils and rock types from the Apollo 15, 16, and 17 lunar sites. Proc. 5th Lunar Sci. Conf. 1087-1096.
- Neal C.R., Taylor L.A., Patchen A.D., Hughes S.S. and Schmitt R.A. (1990a) The significance of fractional crystallization in the petrogenesis of Apollo 17 Type A and B high-Ti basalts. *Geochim. Cosmochim. Acta* 54, 1817-1833.
- Neal C.R. and Taylor L.A. (1993) Catalog of Apollo 17 rocks. Vol. 2 Basalts
- Nord G.L., Lally J.S., Heuer A.H., Christie J.M., Radcliffe S.V., Fisher R.M. and Griggs D.T. (1974) A mineralogical study of rock 70017, an ilmenite-rich basalt, by high voltage electron microscopy (abs). *Lunar Sci.* V, 556-558. Lunar Planetary Institute, Houston
- Nyquist L.E. (1977) Lunar Rb-Sr chronology. *Phys. Chem. Earth* 10, 103-142.
- Nyquist L.E., Bansal B.M. and Wiesmann H. (1975a) Rb-Sr ages and initial ⁸⁷Sr/⁸⁶Sr for Apollo 17 basalts and KREEP basalt 15386. Proc. 6th Lunar Sci. Conf. 1445-1465.
- Nyquist L.E., Bansal B.M. and Wiesmann H. (1976a) Sr isotopic constraints on the petrogenesis of Apollo 17 mare basalts. Proc. 7th Lunar Sci. Conf. 1507-1528.
- Osborne M.D., Parkin K.M. and Bums R.G. (1978) Temperature-dependence of Fe-Ti spectra in the visible region: implications to mapping Ti concentrations of hot planetary surfaces. Proc. 9th Lunar Planet. Sci. Conf. 2949-2960.
- Paces J.B., Nakai S., Neal C.R., Taylor L.A., Halliday A.N. and Lee D.-C. (1991) A strontium and neodymium isotopic study of Apollo 17 high-Ti mare basalts: Resolution of ages, evolution of magmas, and origin of source heterogeneities. *Geochim. Cosmochim. Acta* 55, 2025-2043.
- Papike J.J., Bence A.E. and Lindsley D.H. (1974) Mare basalts from the Taurus-Littrow region of the moon. Proc. 5th Lunar Sci. Conf. 471-504.
- Papike J.J., Hodges F.N., Bence A.E., Cameron M. and Rhodes J.M. (1976) Mare basalts: Crystal chemistry, mineralogy and petrology. *Rev. Geophys. Space Phys.* 14, 475-540.
- Petrowski C., Kerridge J.F. and Kaplan I.R. (1974) Light element geochemistry of the Apollo 17 site. Proc. 5th Lunar Sci. Conf. 1939-1948.
- Philpotts J.A., Schuhmann S., Kouns C.W., Lum R.K.L. and Winzer S. (1974a) Origin of Apollo 17 rocks and soils. Proc. 5th Lunar Sci. Conf. 1255-1267.
- Phinney D., Kahl S.B. and Reynolds J.H. (1975) ⁴⁰Ar-³⁹Ar dating of Apollo 16 and 17 rocks. Proc. 6th Lunar Sci. Conf. 1593-1608.
- Rankenburg K., Brandon A.D. and Neal C.R. (2006) Neodymium isotope evidence for a chondritic composition of the Moon. *Science* 312, 1369-1372.
- Rhodes J.M., Rodgers K.V., Shih C., Bansal B.M., Nyquist L.E., Wiesmann H. and Hubbard N.J. (1974a) The relationships between geology and soil chemistry at the Apollo 17 landing site. Proc. 5th Lunar Sci. Conf. 1097-1117.
- Rhodes J.M., Hubbard N.J., Wiesmann H., Rodgers K.V., Brannon J.C. and Bansal B.M. (1976a) Chemistry, classification, and petrogenesis of Apollo 17 mare basalts. Proc. 7th Lunar Sci. Conf. 1467-1489.
- Roedder E. and Weiblen P.W. (1975a) Anomalous low-K silicate melt inclusions in ilmenite from Apollo 17 basalts. Proc. 6th Lunar Sci. Conf. 147-164.
- Rose H.J., Cuttitta F., Berman S., Brown F.W., Carron M.K., Christian R.P., Dwornik E.J. and Greenland L.P. (1974a)

- Chemical composition of rocks and soils at Taurus-Littrow. Proc. 5th Lunar Sci. Conf. 1119-1133.
- Rose H.J., Brown F.W., Carron M.K., Christian R.P., Cuttitta F., Dwornik E.J. and Ligon D.T. (1974b) Composition of some Apollo 17 samples (abs). Lunar Sci. V, 645-647.
- Rose H.J., Baedeker P.A., Berman S., Christian R.P., Dwornik E.J., Finkelman R.B. and Schnepfe M.M. (1975a) Chemical composition of rocks and soils returned by the Apollo 15, 16, and 17 missions. Proc. 6th Lunar Sci. Conf. 1363-1373.
- Rose H.J., Christian R.P., Dwornik E.J. and Schnepfe M.M. (1975b) Major elemental analysis of some Apollo 15, 16, and 17 samples (abs). Lunar Sci. VI, 686-688. Lunar Planetary Institute, Houston
- Rutherford M.J., Hess P.C. and Daniel G.H. (1974a) Experimental liquid line of descent and liquid immiscibility for basalt 70017. Proc. 5th Lunar Sci. Conf. 569-583.
- Sato M. (1976a) Oxygen fugacity and other thermochemical parameters of Apollo 17 high-Ti basalts and their implications on the reduction mechanism. Proc. 7th Lunar Sci. Conf. 1323-1344.
- Schaeffer O.A. and Husain L. (1973) Isotopic ages of Apollo 17 lunar material (abs). EOS Trans. AGU 54, 614.
- Schaeffer O.A. and Husain L. (1974) Chronology of lunar basin formation. Proc. 5th Lunar Sci. Conf. 1541-1555.
- Schaeffer G.A. and Schaeffer O.A. (1977a) ³⁹Ar/⁴⁰Ar ages of lunar rocks. Proc. 8th Lunar Sci. Conf. 2253-2300.
- Schaeffer G.A. and Schaeffer O.A. (1977b) ³⁹Ar/⁴⁰Ar ages of lunar rocks (abs). Lunar Sci. VIII, 840-842. Lunar Planetary Institute, Houston
- Schwerer F.C. and Nagata T. (1976) Ferromagnetic-superparamagnetic granulometry of lunar surface materials. Proc. 7th Lunar Sci. Conf. 759-778.
- Shih C.-Y., Haskin L.A., Wiesmann H., Bansal B.M. and Brannon J.C. (1975a) On the origin of high-Ti mare basalts. Proc. 6th Lunar Sci. Conf. 1255-1285.
- Shih C.-Y., Wiesmann H. and Haskin L.A. (1975b) On the origin of high-Ti mare basalts (abs). Lunar Sci. VI, 735-737. Lunar Planetary Institute, Houston
- Stephenson A., Collinson D.W. and Runcorn S.K. (1974) Lunar magnetic field paleointensity determinations on Apollo 11, 16, and 17 rocks. Proc. 5th Lunar Sci. Conf. 2859-2871.
- Stephenson A., Runcorn S.K. and Collinson D.W. (1975) On changes in intensity of the ancient lunar magnetic field. Proc. 6th Lunar Sci. Conf. 3049-3062.
- Tilton G.R. and Chen J.H. (1979) Lead isotope systematics of three Apollo 17 mare basalts. Proc. 10th Lunar Planet. Sci. Conf. 259-274.
- Unruh D.M., Stille P., Patchett P.J. and Tatsumoto M. (1984) Lu-Hf and Sm-Nd evolution in lunar mare basalts. Proc. 14th Lunar Planet. Sci. Conf. in J. Geophys. Res. 88, B459-B477.
- Usselman T.M., Lofgren G.E., Donaldson C.H. and Williams R.J. (1975) Experimentally reproduced textures and mineral chemistries of high-titanium mare basalts. Proc. 6th Lunar Sci. Conf. 997-1020.
- Wolfe E.W., Bailey N.G., Lucchitta B.K., Muehlberger W.R., Scott D.H., Sutton R.L and Wilshire H.G. (1981) The geologic investigation of the Taurus-Littrow Valley: Apollo 17 Landing Site. US Geol. Survey Prof. Paper, 1080, pp. 280.
- Yokoyama Y., Reys J.L. and Guichard F. (1974) ²²Na-²⁶Al chronology of lunar surface processes. Proc. 5th Lunar Sci. Conf. 2231-2247.