

**76535**  
Troctolite  
155.5 grams



Figure 1: Photograph of 76535 taken during PET (scale and cube are 1 cm). Photo # S-73-19425

**Introduction**

Troctolite 76535 is without doubt the most interesting sample returned from the Moon! It is a colorful, pristine, coarse-grained, plutonic rock that has had a slow cooling history. It is very old and has not been damaged by shock events. 76535 was collected as part of a rake sample at Station 6. This sample has been widely distributed, but its origin is still debated.

Figure 1 shows the main mass of 76535 before processing. The sample is friable, separating easily at the grain boundaries. Closeup photos of small pieces show the granular texture of the olivine and plagioclase

(figure 2). White plagioclase grains (0.2 to 0.7 mm) are translucent to slightly milky, while lustrous olivine grains (0.2 to 0.3 mm) occur in clusters and are honey-brown in color. Plagioclase shows nice striations on flat cleavage surfaces.

**Petrography**

Gooley et al. (1974) and Dymek et al. (1975) describe lunar sample 76535 as a coarse-grained, olivine-plagioclase cumulate that shows evidence of extensive annealing and re-equilibration (figure 3). This rock has a granular polygonal texture with smooth, curved

**Mineralogical Mode**

	<b>Gooley et al. 1974</b>	<b>Dymek et al. 1975</b>	<b>Warren 1993</b>
Olivine	37 vol. %	35 %	
Plagioclase	58	60	50
Orthopyroxene	4	5	

grain boundaries and abundant 120 degree junctions resulting from the slow process of grain coarsening leading to a mineral fabric with minimum surface area. Stewart (1975) used the grain size of 76535 (2 to 3 mm) and various assumptions to calculate the interval of annealing ( $\sim 10^8$  y) in the temperature range 100 to 600 deg C. Stewart termed this “Apollonian” metamorphism (*although the term did not stick*).

Figure 4 shows that 76535 may be considered as an end member of the Mg suite of lunar highland rocks.

Small patches of vermicular “symplectite” intergrowth of Cr-spinel+high-Ca pyroxene+/-low-Ca pyroxene are a unique feature of this rock and are generally found at grain boundaries between olivine and pyroxene (Gooley et al. 1974; Dymek et al. 1975; Bell et al. 1975). Gooley et al., Dymek et al. and Schwartz and McCallum (1999) have used to reaction relations of this mineral assemblage to calculate the depth of origin of this plutonic rock ( $\sim 47$  km).



Figure 2: Photograph of small piece of 76535 (scale in mm). Note striations on plagioclase. Photo # S-73-19614




Figure 3: Photomicrograph of thin section of 76535 (with partially crossed nicols). Photo is 3 mm across. Photo # S-76-20796.




Figure 4: Plagioclase vs. low-Ca pyroxene plot showing that 76535 is the end-member of the Mg-suite of plutonic rocks. Fields are from James and Flohr (1983) and James et al. (1989).

Small patches of mesostasis (termed “mosaic assemblages”) are found to contain minor minerals including: baddeleyite, low-Ca pyroxene, high-Ca pyroxene, apatite, “whitlockite”, Cr-spinel, metal, K-Ba feldspar, and “pyrochlore” (Gooley et al. 1974; Dymek et al. 1975).

Oriented needles of Fe-Ni-Co metal are found in plagioclase (Bell et al. 1975).

Based on identical mineral chemistry, Warren et al. (1987) apparently found at least two additional pieces of troctolite similar to 76535 in the “coarse fines” from the soil samples (76504,12 and 76034,90).

### Mineralogy

**Olivine:** Olivine is very homogeneous at Fo<sub>87.3</sub> (Dymek et al. 1975; Bersch et al. 1991).

**Pyroxene:** Anhedral grains of orthopyroxene (Wo<sub>0.9</sub>En<sub>84.1</sub>Fs<sub>11.3</sub>) up to 3 mm occur interstitial to olivine and plagioclase (Dymek et al. 1975). Figure 5 shows the composition of mafic minerals in 76535. Hevilon and Crozaz (1989) used the ion microprobe to determine the trace element content of pyroxene. Careful crystal structure refinement of single crystal X-ray study of orthopyroxene indicate closure temperature for cation ordering at about 550 deg C (Domeneghetti et al. 2001). Smyth (1975) also studied the crystal structure of opx from 76535.




Figure 5: Pyroxene and olivine composition of 76535. Data replotted from Dymek et al. (1975) and Gooley et al. (1974).

**Plagioclase:** Plagioclase is homogeneous at An<sub>96.2</sub> (Dymek et al. 1975). White to translucent plagioclase grains show striations due to twinning (LSPET 1973; Phinney et al. 1974; Gooley 1974). Smyth (1986) performed crystal structure refinement of anorthite using plagioclase from 76535 to determine the position of the cations in the structure. Nord (1976) studied the inclusions found in plagioclase. The trace element content of plagioclase has been reported by Haskin et al. (1975), Hansen et al. (1979), Steele et al. (1980), and Hevilon and Crozaz (1989).

**Augite:** Small grains of augite are found in symplectite and mesostasis areas. Dymek et al. (1975) report Wo<sub>43.6</sub>En<sub>48.5</sub>Fs<sub>4.4</sub>

**Spinel:** Dymek et al. (1975) and Haggerty (1975) gives the composition of Cr-spinel, which is found intergrown with augite in symplectite.

**Baddeleyite:** Relatively large grains of ZrO<sub>2</sub> are present.

**Metal:** Gooley et al. (1974) and Ryder et al. (1980) report the composition of metal grains in 76535.

**Phosphate:** The composition of both apatite (F 2.95% and Cl 1.18%) and whitlockite (high REE) are given in Dymek et al. (1975).

**Troilite:** Small grains have been reported.

**Pyrochlore:** Reported by Dymek et al. (1975).




Figure 6: Normalized rare earth element diagram for lunar troctolite 76535. Data from Haskin et al. (1975).




Figure 8: Rb-Sr isochron diagram for 76535. From Papanastassiou and Wasserburg (1976).




Figure 7: U-Pb concordia diagram for 76535. From Premo and Tatsumoto (1992).




Figure 9: Sm-Nd isochron for 76535 troctolite. From Lugmair et al. (1976).

## Chemistry

Rhodes et al. (1974), Haskin et al. (1974) and Wiesmann and Hubbard (1975) reported the bulk chemical composition of 76535 (table 1 and figure 6). Morgan et al. (1974) and Wolf et al. (1979) reported the trace siderophile and volatile elements.

Haskin et al. used the whole-rock composition and known distribution coefficients to calculate the probable parent liquid. They concluded that the rock may have had ~16% trapped liquid when it originally crystallized from the melt.

### Summary of age data for 76535

4.19 ± 0.02	K-Ar	Husain and Schaeffer (1975)
4.16 ± 0.04	K-Ar	Huneke and Wasserburg (1975)
4.27 ± 0.08	K-Ar	Bogard et al. (1975)
4.61 ± 0.07	Rb-Sr	Papanastassiou and Wasserburg (1976)
4.26 ± 0.02	Sm-Nd	Lugmair et al. (1976)
4.330 ± 0.064	Sm-Nd	Premo and Tatsumoto (1992)
4.27 ± 0.03	Pb-Pb	Hinthorne et al. (1975)
4.236 ± 0.015	U-Pb	Premo and Tatsumoto (1992)

## Radiogenic age dating

Heroic efforts have been made to age date troctolite 76535 and the data from different decay schemes seem to be at odds (see Summary). Most recently, Premo and Tatsumoto (1992) have carefully considered the age of 76535 and concluded that it was formed between 4.23 and 4.26 b.y. (figure 7).

**Table 1. Chemical composition of 76535.**

reference weight	Rhodes 74	Haskin 74	Wiesmann 75	Morgan 74	Kieth 74
	1.3 g			151 mg	155 g
SiO <sub>2</sub> %	42.88 (a)				
TiO <sub>2</sub>	0.05 (a)				
Al <sub>2</sub> O <sub>3</sub>	20.73 (a)				
FeO	4.99 (a)				
MnO	0.07 (a)				
MgO	19.09 (a)				
CaO	11.41 (a)				
Na <sub>2</sub> O	0.23 (d)				
K <sub>2</sub> O	0.03 (a)	0.028 (c)			0.0283 (f)
P <sub>2</sub> O <sub>5</sub>	0.03 (a)				
S %	0 (a)				
sum					
Sc ppm					
V					
Cr	753 (a)				
Co					
Ni	25 (a)			44 (e)	
Cu					
Zn	1			1.2 (e)	
Ga					
Ge				0.0017 (e)	
As					
Se				0.0041 (e)	
Rb	0.2	0.24 (c)		0.2 (e)	
Sr	111	114 (c)			
Y	4.4				
Zr	12	23.6 (c)			
Nb	1.2				
Mo					
Ru					
Rh					
Pd ppb					
Ag ppb				0.12 (e)	
Cd ppb					
In ppb					
Sn ppb					
Sb ppb				0.014 (e)	
Te ppb				0.28 (e)	
Cs ppm				0.014 (e)	
Ba		33 (c)			
La		1.51 (c)			
Ce		3.8 (c)			
Pr					
Nd		2.3 (c)			
Sm		0.61 (c)			
Eu		0.73 (c)			
Gd		0.73 (c)			
Tb					
Dy		0.8 (c)			
Ho					
Er		0.53 (c)			
Tm					
Yb		0.56 (c)			
Lu		0.079 (c)			
Hf		0.52 (c)			
Ta					
W ppb					
Re ppb				0.0012 (e)	
Os ppb					
Ir ppb				0.0054 (e)	
Pt ppb					
Au ppb				0.0025 (e)	
Th ppm					0.19 (f)
U ppm				0.0194 (e)	0.054 (f)

technique (a) XRF, (b) INAA, (c) IDMS, (d) AA, (e) RNAA, (f) counting




Figure 10: Reflectance spectra of Apollo 17 rocks including troctolite 76535. From Charette and Adams (1977).

The various age dating studies of troctolite 76535 may provide an interesting study in the preservation of radiogenic information through the course of major metamorphic change. Careful study of Rb-Sr, Sm-Nd, U-Pb and  $^{40}\text{Ar}/^{39}\text{Ar}$  systematics has yielded a broad range of apparent isotopic closure ages; 4.61 to 4.16 b.y. (see Summary). The Rb-Sr isochron (figure 8) is based on Rb-rich inclusions in olivine, whereas the Sm-Nd isochron (figure 9) is based on REE included in pyroxene, plagioclase and accessory phases, exclusive of olivine.

Domeneghetti et al. (2001) and Nord (1976) note that this apparent age discrepancy is consistent with what is known about cation ordering in orthopyroxene and subsolidus exsolution of inclusions found in plagioclase. The Rb-Sr isochron presumably dates the isolation of Rb-rich inclusions in olivine (?) and is apparently insensitive to the metamorphism that produced the texture of the rock, while the Sm-Nd, U-Pb and  $^{40}\text{Ar}/^{39}\text{Ar}$  ages involve lower temperature mineral phases that are more sensitive to subsequent metamorphism and closure to movement of radiogenic elements at a later time. The study of rare gases by Chaffee et al. (1981) also shows that the different minerals in 76535 have different, mineral-specific, isotopic closure ages.




Figure 11: Relative abundance of trapped rare gases in 76535. From Chaffee et al. (1981).




Figure 12: Xenon isotopic data for olivine and plagioclase separates from 76535. From Chaffee et al. (1981).

### Cosmogenic isotopes and exposure ages

Bogard et al. (1975), Crozaz et al. (1974) and Lugmair et al. (1976) reported cosmic ray exposure ages of  $195 \pm 10$  m.y.,  $211 \pm 7$  m.y., and  $233 \pm 16$  m.y., respectively.

Premo and Tatsumoto (1992) show a hint of a lower intercept age at  $62 \pm 32$  m.y., suggesting Pb disturbance at the time of excavation.

Kieth et al. (1974) reported the cosmic ray induced activity of 76535.

## Other Studies

Charette and Adams (1977) have recorded the spectral reflectance of 76535 (figure 10). Braddy et al. (1975) reported a fission track age of 4.07 b.y.

Hohenberg et al. (1980) and Chaffee et al. (1981) have carefully studied “excess” fission xenon and trapped solar wind noble gases in troctolite 76535 (figures 11, 12). Stepwise heating of separated olivine and plagioclase showed evidence for *in-situ* decay of <sup>244</sup>Pu leading to fission Xe ages of 4.50 b.y. and 4.25 b.y. respectively (consistent with Rb-Sr and Sm-Nd ages above).


## Photo Numbers

S73-19425  
 S73-19458  
 S73-19459  
 S73-19601  
 S73-19614  
 S73-19622  
 S73-20409B  
 S76-20796

## Processing

Two 2 gram pieces of 76535 were used by Rhodes et al. (1994) and Haskin et al. (1994) to obtain the chemical composition.

The sample is very friable and broke up during initial sawing. There are probably other pieces of this sample in the residue of the collection bag (76530).



## References for 76535

- Bell P.M. and Mao H.K. (1975) Cataclastic plutonites: Possible keys to the evolutionary history of the early Moon (abs). *Lunar Sci.* **VI**, 34-35. Lunar Planetary Institute, Houston.
- Bell P.M., Mao H.K., Roedder E. and Weiblen P.W. (1975) The problem of the origin of symplectites in olivine-bearing lunar rocks. *Proc. 6<sup>th</sup> Lunar Sci. Conf.* 231-248.
- Bersch M.G., Taylor G.J., Keil K. and Norman M.D. (1991) Mineral compositions in pristine lunar highland rocks and the diversity of highland magmatism. *Geophys. Res. Lett.* **18**, 2085-2088.
- Bogard D.D. and Nyquist L.E. (1974) 76535: An old lunar rock? (abs) *Lunar Sci.* **V**, 70-72. Lunar Planetary Institute, Houston.
- Bogard D.D., Nyquist L.E., Bansal B.M., Wiesmann H. and Shih C.-Y. (1975) 76535: An old lunar rock. *Earth Planet. Sci. Lett.* **26**, 69-80.
- Braddy D., Hutcheon I.D. and Price P.B. (1975a) Crystal chemistry of Pu and U and concordant fission track ages of lunar zircons and whitlockites. *Proc. 6<sup>th</sup> Lunar Sci. Conf.* 3581-3600.
- Braddy D., Hutcheon I.D. and Price P.B. (1975b) Crystal chemistry of Pu and U and concordant fission track ages of lunar zircons and whitlockites (abs). *Lunar Sci.* **VI**, 77-79. Lunar Planetary Institute, Houston.
- Butler P. (1973) **Lunar Sample Information Catalog Apollo 17**. Lunar Receiving Laboratory. MSC 03211 Curator's Catalog. pp. 447.
- Charette M.P. and Adams J.B. (1977) Spectral reflectance of lunar highland rocks (abs). *Lunar Sci.* **VIII**, 172-174. Lunar Planetary Institute, Houston.
- Caffee M., Hohenberg C. and Hudson B. (1981a) Troctolite 76535: A study in the preservation of early isotopic records (abs). *Lunar Planet. Sci.* **XII**, 120-122. Lunar Planetary Institute, Houston.
- Caffee M., Hohenberg C.M. and Hudson B. (1981b) Troctolite 76535: A study in the preservation of early isotopic records. *Proc. 12<sup>th</sup> Lunar Planet. Sci. Conf.* 99-115.
- Crozaz G., Drozd R., Hohenberg C., Morgan C., Ralston C., Walker R. and Yuhas D. (1974a) Lunar surface dynamics: Some general conclusions and new results from Apollo 16 and 17. *Proc. 5<sup>th</sup> Lunar Sci. Conf.* 2475-2499.
- Domeneghetti M.C., McCallum I.S., Schwartz J.M., Camara F., Zema M., McCammon C. and Ganguly J. (2001) Complex cooling histories of lunar troctolite 76535 and Stillwater orthopyroxenite SC-936 (abs#1151). *Lunar Planet. Sci.* **XXX**, Lunar Planetary Institute, Houston.
- Dymek R.F., Albee A.L. and Chodos A.A. (1975b) Comparative petrology of lunar cumulate rocks of possible primary origin: Dunite 72415, troctolite 76535, norite 78235, and anorthosite 62237. *Proc. 6<sup>th</sup> Lunar Sci. Conf.* 301-341.
- Gooley R.C., Brett R., Warner J.L. and Smyth J.R. (1974) A lunar rock of deep crustal origin: Sample 76535. *Geochim. Cosmochim. Acta* **38**, 1329-1339.
- Hansen E.C., Steele I.M. and Smith J.V. (1979a) Lunar highland rocks: Element partitioning among minerals 1: Electron microprobe analyses of Na, K, and Fe in plagioclase; mg partitioning with orthopyroxene. *Proc. 10<sup>th</sup> Lunar Planet. Sci. Conf.* 627-638.
- Haskin L.A., Shih C.-Y., Bansal B.M., Rhodes J.M., Wiesmann H. and Nyquist L.E. (1974a) Chemical evidence for the origin of 76535 as a cumulate. *Proc. 5<sup>th</sup> Lunar Sci. Conf.* 1213-1225.
- Haskin L.A., Shih C.-Y., Bansal B.M., Rhodes J.M., Wiesmann H. and Nyquist L.E. (1974b) Chemical evidence for the origin of 76535 as a cumulate (abs). *Lunar Sci.* **V**, 313-315. Lunar Planetary Institute, Houston.
- Hinthorne J.R., Conrad R.L. and Andersen C.A. (1975) Lead-lead and trace element abundances in lunar troctolite, 76535 (abs). *Lunar Sci.* **VI**, 373-375. Lunar Planetary Institute, Houston.
- Hinthorne J.R., Andersen C.A., Conrad R.L. and Lovering J.F. (1979) Single-grain <sup>207</sup>Pb/<sup>206</sup>Pb and U/Th age determinations with a 10-micron spatial resolution using the ion microprobe mass analyzer (IMMA). *Chem. Geology* **25**, 271-303.
- Heavilon C.F. and Crozaz G. (1989) REE and selected minor and trace element microdistributions in some pristine lunar highlands rocks (abs). *Lunar Planet. Sci.* **XX**, 398-399. Lunar Planetary Institute, Houston.
- Hohenberg C.M., Hudson B., Kennedy B.M. and Podosek F.A. (1980) Fission xenon in troctolite 76535. In **Proc. Conf. Lunar Highlands Crust**. *Geochim. Cosmochim. Acta*, Suppl. 12. Pergamon Press. 419-439. Lunar Planetary Institute, Houston.
- Huneke J.C. and Wasserburg G.J. (1975) Trapped <sup>40</sup>Ar in troctolite 76535 and evidence for enhanced <sup>40</sup>Ar-<sup>39</sup>Ar age



- plateaus (abs). *Lunar Sci.* **VI**, 417-419. Lunar Planetary Institute, Houston.
- Husain L. and Schaeffer O.A. (1975) Lunar evolution: The first 600 million years. *Geophys. Res. Lett.* **2**, 29-32.
- James O.B. and Flohr M.K. (1983) Subdivision of the Mg-suite noritic rocks into Mg-gabbro-norites and Mg-norites. *Proc. 13<sup>th</sup> Lunar Planet. Sci. Conf. in J. Geophys. Res.*, A603-A614.
- James O.B. (1982) Subdivision of the Mg-suite plutonic rocks into Mg-norites and Mg-gabbro-norites (abs). *Lunar Planet. Sci.* **XIII**, 360-362. Lunar Planetary Institute, Houston.
- Keith J.E., Clark R.S. and Bennett L.J. (1974a) Determination of natural and cosmic ray induced radionuclides in Apollo 17 lunar samples. *Proc. 5<sup>th</sup> Lunar Sci. Conf.* 2121-2138.
- LSPET (1973) Apollo 17 lunar samples: Chemical and petrographic description. *Science* **182**, 659-672.
- LSPET (1973) Preliminary Examination of lunar samples. Apollo 17 Preliminary Science Rpt. NASA SP-330. 7-1 – 7-46.
- Lugmair G.W., Marti K., Kurtz J.P. and Scheinin N.B. (1976a) History and genesis of lunar troctolite 76535 or: How old is old? *Proc. 7<sup>th</sup> Lunar Sci. Conf.* 2009-203.
- Lugmair G. and Marti K. (1978) Lunar initial <sup>143</sup>Nd/<sup>144</sup>Nd: differential evolution of the lunar crust and mantle. *Earth Planet. Sci. Lett.* **39**, 349-357.
- Meyer C. (1994) Catalog of Apollo 17 rocks: Volume 4. Curator's Office JSC 26088 pp. 644
- Morgan J.W., Ganapathy R., Higuchi H., Krahenbuhl U. and Anders E (1974a) Lunar basins: Tentative characterization of projectiles, from meteoritic dements in Apollo 17 boulders. *Proc. 5<sup>th</sup> Lunar Sci. Conf.* 1703-1736.
- Muehlberger et al. (1973) Documentation and environment of the Apollo 17 samples: A preliminary report. *Astrogeology* **71** 322 pp superceeded by *Astrogeology* **73** (1975) and by Wolfe et al. (1981)
- Muehlberger W.R. and many others (1973) Preliminary Geological Investigation of the Apollo 17 Landing Site. *In Apollo 17 Preliminary Science Report.* NASA SP-330.
- Nord G.L., Ross M. and Huebner J.S. (1976) Lunar troctolite 76535: Mineralogical investigations (abs). *Lunar Sci.* **VII**, 628-630. Lunar Planetary Institute, Houston
- Nord G.L., Heubner J.S. and Ross M. (1977) Structure, composition, and significance of "G-P" zones in 76535 orthopyroxene (abs). *Lunar Planet. Sci.* **VIII**, 732-734. Lunar Planetary Institute, Houston.
- Papanastassiou D.A. and Wasserburg G.J. (1976a) Rb-Sr age of troctolite 76535. *Proc. 7<sup>th</sup> Lunar Sci. Conf.* 2035-2054.
- Phinney W.C., Simonds C.H. and Warner J. (1974) Description, Classification and Inventory of Apollo 17 Rake Samples from Station 6. Curator's Catalog, pp. 46.
- Premo W.R. and Tatsumoto M. (1992a) U-Th-Pb, Rb-Sr, and Sm-Nd isotopic systematics of lunar troctolite cumulate 76535: Implications on the age and origin of this early lunar, deep-seated cumulate. *Proc. 22<sup>nd</sup> Lunar Planet. Sci. Conf.* 381-397. Lunar Planetary Institute, Houston.
- Rhodes J.M., Rodgers K.V., Shih C., Bansal B.M., Nyquist L.E., Wiesmann H. and Hubbard N.J. (1974a) The relationships between geology and soil chemistry at the Apollo 17 landing site. *Proc. 5<sup>th</sup> Lunar Sci. Conf.* 1097-1117.
- Ryder G. and Norman M.D. (1979a) Catalog of pristine non-mare materials Part 1. Non-anorthosites, revised. NASA-JSC Curatorial Facility Publ. JSC 14565, Houston. 147 pp.
- Ryder G., Norman M.D. and Score R.A. (1980a) The distinction of pristine from meteorite-contaminated highlands rocks using metal compositions. *Proc. 11<sup>th</sup> Lunar Planet. Sci. Conf.* 471-479.
- Smyth J.R. (1975) Intracrystalline cation order in a lunar crustal troctolite. *Proc. 6<sup>th</sup> Lunar Sci. Conf.* 821-832.
- Smyth J.R. (1986) Crystal structure refinement of a lunar anorthite, An<sub>94</sub>. *Proc. 17<sup>th</sup> Lunar Planet. Sci. Conf.* in *J. Geophys. Res.* **91**, E91-97.
- Steele I.M., Hutcheon I.D. and Smith J.V. (1980) Ion microprobe analysis and petrogenetic interpretations of Li, Mg, Ti, K, Sr, Ba in lunar plagioclase. *Proc. 11<sup>th</sup> Lunar Planet. Sci. Conf.* 571-590.
- Stewart D.B. (1975) Apollonian metamorphic rocks—The products of prolonged subsolidus equilibration (abs). *Lunar Sci.* **VI**, 774-776. Lunar Planetary Institute, Houston.
- Schwartz J.M. and McCallum I.S. (1999) Inferred depths of formation of spinel cataclasites and troctolitic granulite, 76535 using new thermodynamic data for Cr-spinel (abs1308). *Lunar Planet. Sci.* **XXX**, Lunar Planetary Institute, Houston.

Tera F. and Wasserburg G.J. (1974) U-Th-Pb systematics on lunar rock: and inferences about lunar evolution and the age of the Moon. *Proc. 5<sup>th</sup> Lunar Sci. Conf.* 1571-1599.

Tera F., Papanastassiou D.A. and Wasserburg G.J. (1974a) Isotopic evidence for a terminal lunar cataclysm. *Earth Planet. Sci. Lett.* **22**, 1-21.

Warren P.H. and Wasson J.T. (1977) Pristine nonmare rocks and the nature of the lunar crust. *Proc. 8<sup>th</sup> Lunar Sci. Conf.* 2215-2235.

Warren P.H. (1993) A concise compilation of petrologic information on possibly pristine nonmare Moon rocks. *Am. Mineral.* **78**, 360-376.

Wiesmann H. and Hubbard N.J. (1975) A compilation of the Lunar Sample Data Generated by the Gast, Nyquist and Hubbard Lunar Sample PI-Ships. Unpublished. JSC

Wolf R., Woodrow A. and Anders E. (1979) Lunar basalts and pristine highland rocks: Comparison of siderophile and volatile elements. *Proc. 10<sup>th</sup> Lunar Planet. Sci. Conf.* 2107-2130.

Wolfe E.W., Bailey N.G., Lucchitta B.K., Muehlberger W.R., Scott D.H., Sutton R.L. and Wilshire H.G. (1981) The geologic investigation of the Taurus-Littrow Valley: Apollo 17 Landing Site. US Geol. Survey Prof. Paper, 1080, pp. 280.