

77215
Cataclastic Norite
846.4 grams



Figure 1: Tray full of pieces of 77215. Some pieces are covered with “off-white” patina including obvious zap pits. NASA photo # S73-17779.

Introduction

Apollo 17 sample 77215 represents the oldest lithology in the Station 7 boulder. It is cut by a dark dike and surrounded by two generations of melt rock (see figure 2 in section on 77075). 77215 is probably a brecciated pigeonite-anorthite cumulate (figure 3), but the small size of the remnant lithic clasts prevents any certain determination of cumulate origin (Chao et al. 1976). The pigeonite has inverted to orthopyroxene with augite exsolution. The presence of undevitrified noritic glass in 77215 is significant to understanding the thermal history of this boulder.

77215 was collected from the obvious large white clast in the Station 7 boulder (Wolfe and others 1981). While it appeared “off-white” or “light-gray” in the surface photography, the fresh surfaces of the sample proved

to be pure white in the laboratory (figure 1). One piece of 77215 (piece 19) includes the dark dike material of 77075 (figure 4). Other pieces contain thin dark veins (as in sample 77077).

This sample and others from the Station 7 boulder were studied by the International Consortium led by Ed Chao (see summaries by Chao et al. 1976, Winzer et al. 1977 and Minkin et al. 1978). The results on 77215 were also summarized in the catalog by Meyer (1994).

The Consortium was unable to provide exactly similar subsamples to individual member for study and analysis, because of the clastic nature of the 77215. Subsamples were assigned to individual consortium participants on the basis of suitability for their



Figure 2: Sawn surface of 77215,16 showing relict igneous texture. NASA photo # S83-34595. One cm cube for scale.




Figure 3: Photomicrograph of a norite clast in thin section 77215,139 (from Chao et al. 1976). Small inclusions of K-feldspar(K) are characteristic.

Mineralogical Mode of 77215

	Chao et al. 1974	Chao et al. 1976
Plagioclase	52 vol. %	54
Orthopyroxene	45	41.5
Augite		0.4
Silica		1.8
Troilite		0.6
Ilmenite		0.1
Metal		0.3
Spinel		0.2
K-feldspar		0.3
Glass		1.0



Figure 4: Slab 77215,82 showing dark dike in 77215,19. Note rounded xenocrysts in fine-grained dike. Cube is 1 cm for scale. NASA photo # S75-21979.

experiments and the resulting data cannot now be exactly correlated (Minkin et al. 1978).

(see the transcript of the astronauts, provided in the section on 77075)

Petrography

77215 is a friable breccia that broke up into many pieces on the way back from the moon (figure 1). Some pieces have small areas of unbrecciated norite with primary igneous texture (figures 2 and 3). However, most of the lithic clasts in 77215 have been intensely granulated or smeared out to form schlieren, so that the relict host rock(s) are only represented by very small clasts (Chao et al. 1976).

The modal mineralogy of 77215 is approximately 41% orthopyroxene and 54% plagioclase with trace amounts of troilite, ilmenite, clinopyroxene, spinel, silica, K-feldspar, and other trace phases. Although it contains some gray glass, it is low in Ir and Ni. Thus, 77215 is (or rather was) a pristine norite that has been shocked and crushed in place. Most of the plagioclase has not been converted to maskelynite by the mild shock pressure.

The plagioclase and pyroxene in 77215 have a narrow range of chemical composition (An_{90-91} and $Wo_{3-5}En_{63-68}Fs_{29-32}$) and plot in the range of lunar norite (figure 7). Pyroxenes in 77215 show some of the features of “inverted pigeonites”. Huebner et al. (1975) explain that the misoriented nature of the augite, relative to the orthopyroxene host, is a common feature of



Figure 5: Sawn surfaces of 77215,92 with cm scale. Note obvious clasts in fine-grained, crushed matrix. NASA # S75-21980.




Figure 6: Photomicrograph of thin section of 77215,14. a) plane polarized light, b) cross-polarized light, showing glass matrix. NASA photo #s S79-27400 and 27401. Field of view is 1.4 mm.

pyroxenes that originally crystallized as homogeneous pigeonite crystals at high temperatures. According to Huebner et al., coarse pyroxene exsolution lamellae can form in geologically short periods of time (<30,000 yr.) at elevated temperature. Huebner et al. argue that such conditions could have been met in the upper levels of the lunar crust during early lunar history as a consequence of the initially hot crust of the moon. According to Huebner et al., the exsolved pyroxenes seen in 77215 do not necessarily suggest the deep-seated origin, as originally proposed by Chao et al. (1974).

Significant Clasts

Huebner et al. (1975) report a lithic clast, 3.5 mm long, in thin section 77215,13 that consists entirely of orthopyroxene and plagioclase, in roughly equal proportions. The grain size is 0.5 to 2 mm, and the texture is subophitic. They consider that the clast is an unbrecciated piece of the parental cumulate rock that has escaped granulation.

A gray impact glass clast in 77215,29 was subdivided for detailed study (Chao et al. 1976). The chemical analysis (labeled ,130) of this clast and microprobe analyses (Chao et al. 1976) showed it to have the bulk composition of the norite (figure 8).

Mineralogy

Pyroxene: The pyroxene in 77215 is orthopyroxene (host) with exsolved augite lamellae and blebs (Huebner et al. 1975)(figure 8). Winzer et al. (1977) analyzed pyroxene separates for 77215 and Papike et al. (1994) used the ion microprobe method to determine




Figure 7: Composition of plagioclase and pyroxene in 77215.

the trace element content of orthopyroxene. Anderson and Lindsley (1982) have carefully calculated the equilibrium temperature of pyroxene pairs in 77215 ($T = 850 \pm 30$ deg C).

Plagioclase: Huebner et al. (1975) report that the plagioclase in 77215 is $Or_{0.7}Ab_{8.2}An_{91.1}$, with low Mg and Fe content, typical of plagioclase from plutonic lunar rocks. Chao et al. (1976) report the composition as $An_{88-92}Ab_{11-7}Or_1$. Plagioclase has not been converted to maskelynite by shock. Winzer et al. (1977) provide trace element analysis of plagioclase separates.

K-feldspar: Plagioclase grains frequently contain square to rectangular inclusions of K-feldspar ($Or_{97}Ab_1An_2$) or granitic glass (Chao et al. 1976).

Other: Chao et al. (1976) report “clusters” of accessory minerals consisting of Fe-Co metal, troilite, ilmenite, chromite, anorthite, orthopyroxene, silica, rare augite and whitlockite. Chao et al. (1976) and Huebner et al. (1975) also report grains of Zr-Ti-Ca-Fe oxide.

Glass: Gray glass clasts and veins with the approximate composition of the bulk norite are present in 77215 (Chao et al. 1976). An analysis of this glass is found in table 1.

Chemistry

The consortium created two large, homogeneous splits for accurate chemical analysis (split 77215,45 weighed 1.13 grams and split ,152 weighed 0.62 grams). Table




Figure 8: Composition of orthopyroxene with exsolved augite in 77215 (from Huebner et al. 1975). Rare olivine is out of equilibrium.




Figure 9: Normalized rare-earth-element composition of 77215 and dike (data from Winzer et al. 1977).

1 and figure 9 give the composition of the white norite and the dark dike material. The gray glass has the same composition as the white norite.

The white noritic portion of 77215 is pristine (low Ir, Higuchi and Morgan 1975, Ebihara et al. 1991, Wolf et al. 1979, Warren 1993). The dark dike material has the same composition as 77075.

Radiogenic age dating

The argon 39/40 plateau age for 77215 is significantly lower than the age determined by Rb/Sr or Sm/Nd internal mineral isochrons (as was also the case for norite sample 78235). The best Ar age (3.98 b.y.) seems to be from the plagioclase separate (figure 12), which probably represents the time of formation of the melt sheet (figure 10). The original crystallization age of the parental norite appears to be ~ 4.4 b.y. (Nakamura et al. 1976). This is only one of a few samples of the




Figure 10: Summary diagram for ages determined by Argon release from samples of Station 7 boulder (from Stettler et al. 1978).

original crust of the moon that have been successfully dated (figure 13 and 14).

U-Th-Pb data for 77215 are disturbed (Nunes et al. 1974).

Cosmogenic isotopes and exposure ages

Stettler et al. (1974) determined an Ar exposure age of 27.2 m.y.

Processing

The International Consortium selected subsamples 77215,19,22,29,37,45 and ,58 for study because they are representative of the clast assemblages of this breccia. Some notes on the distribution of these subsamples is given in the appendix to Chao et al. (1976). Detailed description of the splits is recorded in Open File Report 78-511. Butler and Dealing (1974) outlined the original processing and distribution of samples from this boulder. There are 26 thin sections.

List of Photo #s for 77215

S73-17778-17779	tray full
S83-34595	,16 sawn surface
S75-21979	
S75-21992	,19 sawn
S75-21980	,29 sawn

Summary of Age Data for 77215

	Ar-Ar	Rb/Sr	Sm/Nd
Stettler et al. 1974	3.96 ± 0.03 b.y.		
Stettler et al. 1978	3.98 ± 0.03 (plag.)		
Nakamura et al. 1976		4.42 ± 0.04	4.37 ± 0.07

Caution: These ages use old decay constants.




Figure 11: Argon release diagram for 77075 and 77215 (from Stettler et al. 1974).




Figure 12: Ar release diagram for plagioclase in 77215 (from Stettler et al. 1978).





Figure 12: Rb/Sr isochron for 77215 as determined by Nakamura et al. (1976).

Figure 13: Sm-Nd internal mineral isochron for 77215 by Nakamura et al. (1976).

77215 (pieces)




Table 1. Chemical composition of 77215

reference	norite Winzer 74	matrix Winzer 77	glass Winzer 77	black dike	dike	dike		Higuchi 75 ,37	Ebihara 92 ,35		77077 white Warren 78 for comparison
<i>weight</i>	98 mg	,152	,130	,115	,119	,121					
SiO2 %	51.3	51.1	51.1	46.8	47.2	46	(b)				50.9 (e)
TiO2	0.32	0.3	0.37	1.37	1.35	1.32	(c)				0.3 (e)
Al2O3	15.06	13.98	14.32	17.44	16.89	17.75	(b)				16.16(e)
FeO	10.07	10.38	10.32	9.39	9.36	9.04	(b)				8.74 (e)
MnO	0.16	0.17	0.17	0.12	0.12	0.11	(b)				0.15 (e)
MgO	12.56	14.31	13.23	13.16	12.93	12.74	(b)				10.6 (e)
CaO	8.96	8.65	9.08	10.88	10.76	10.94	(b)				9.94 (e)
Na2O	0.43	0.39	0.55	0.65	0.68	0.68	(b)				0.44 (e)
K2O	0.14	0.18	0.15	0.24	0.23	0.24	(b)				0.22 (e)
P2O5	0.11	0.14	0.1	0.28	0.27	0.26	(c)				
S % <i>sum</i>											
Sc ppm											13.8 (e)
V											
Cr	2189	2463	2463	1300	1368	958	(b)				
Co											25.2 (e)
Ni								<3	50	(d)	< 1.7 (e)
Cu											
Zn								3	2.95	(d)	2.84 (e)
Ga											5 (e)
Ge ppb								14.3	47.1	(d)	18.7 (e)
As											
Se								77	83.2	(d)	
Rb	3.54	3.21		6.51	6.48	6.26	(a)	4.9	12.3	(d)	
Sr	105	102	103	171	169	174	(a)				
Y											
Zr	171		147	419			(a)				150 (e)
Nb											
Mo											
Ru											
Rh											
Pd ppb									1.45	(d)	
Ag ppb								0.62	1.89	(d)	
Cd ppb								4.4	4.39	(d)	
In ppb									<0.1	(d)	
Sn ppb											
Sb ppb								0.121	1.04	(d)	
Te ppb								1	1.92	(d)	
Cs ppm								0.18	0.393	(d)	
Ba	166	154	154	350	349	336	(a)				220 (e)
La											9.9 (e)
Ce	27.2	24.6	29.6	84.4	73.3	79.1	(a)				25 (e)
Pr											
Nd	16.8	15.5	18	51.9	51.7	50.8	(a)				16 (e)
Sm	4.68	4.4	5.05	14.4	14.5	13.8	(a)				4.28 (e)
Eu	1.08	1.03	1.01	1.93	1.9	1.97	(a)				1.12 (e)
Gd	6.64	5.21					(a)				
Tb											1 (e)
Dy	7.08	6.64	7.31	19.6	19.4	18.4	(a)				
Ho											
Er	4.51	4.57	4.44	10		10.7	(a)				
Tm											
Yb	4.98	4.88	4.45	8.59	10.5	9.94	(a)				4.5 (e)
Lu	0.766	0.592	0.835	1.76		1.68	(a)				0.67 (e)
Hf						3	(a)				3.4 (e)
Ta											0.38 (e)
W ppb											
Re ppb								0.0047	0.173	(d)	
Os ppb									3.04	(d)	
Ir ppb								0.0221	2.66	(d)	0.0029(e)
Pt ppb											
Au ppb								0.0108	0.557	(d)	0.056(e)
Th ppm											2 (e)
U ppm								0.92	0.799	(d)	0.59 (e)

technique (a) IDMS, (b) AA, (c) colorimetry, (d) RNAA, (e) INAA

Table 2: Composition of 77215

	U ppm	Th ppm	K2O %	Rb ppm	Sr ppm	Nd ppm	Sm ppm	technique
Winzer et al. 1974			0.14	3.54	105	16.8	4.68	IDMS
			0.18	3.21	102	15.5	4.4	IDMS
Nakamura et al. 1976			0.127	6.177	65.46	14.84	4.372	IDMS
			0.0842	2.326	86.81			
Nunes et al. 1974	0.5068	1.993						IDMS
Higuchi et al. 1975	0.92			4.9				RNAA
Ebihara et al. 1992	0.799			12.3				RNAA

References for 77215

- Andersen D.J. and Lindsley D.H. (1982) Application of a two-pyroxene thermometer (abs). *Lunar Planet. Sci. XIII*, 15-16. Lunar Planetary Institute, Houston.
- Bersch M.G., Taylor G.J., Keil K. and Norman M.D. (1991) Mineral compositions in pristine lunar highland rocks and the diversity of highland magmatism. *Geophys. Res. Lett.* **18**, 2085-2088.
- Butler P. (1973) **Lunar Sample Information Catalog Apollo 17**. Lunar Receiving Laboratory. MSC 03211 Curator's Catalog. pp. 447.
- Butler P. and Dealing T.E. (1974) The dissection and consortium allocation of Apollo 17 lunar rocks from the boulder at Station 7. *Earth Planet. Sci. Lett.* **23**, 429-434.
- Chao E.C.T., Minkin J.A. and Thompson C.L. (1974) Preliminary petrographic description and geologic implications of the Apollo 17 Station 7 Boulder Consortium samples. *Earth Planet. Sci. Lett.* **23**, 413-428.
- Chao E.C.T., Minkin J.A. and Thompson C.L. (1976a) The petrology of 77215, a noritic impact breccia. *Proc. 7th Lunar Sci. Conf.* 2287-2308.
- Chao E.C.T., Minkin J.A. and Thompson C.L. (1976b) The petrology of 77215, a noritic impact ejecta breccia (abs). *Lunar Sci. VII*, 129-131. Lunar Planetary Institute, Houston.
- Ebihara M., Wolf R., Warren P.H. and Anders E. (1992) Trace elements in 59 mostly highland moon rocks. *Proc. 22nd Lunar Planet. Sci. Conf.* 417-426. Lunar Planetary Institute, Houston
- Higuchi H. and Morgan J.W. (1975a) Ancient meteoritic component in Apollo 17 boulders. *Proc. 6th Lunar Sci. Conf.* 1625-1651.
- Higuchi H. and Morgan J.W. (1975b) Ancient meteoritic component in Apollo 17 boulders (abs). *Lunar Sci. VI*, 364-366. Lunar Planetary Institute, Houston.
- Huebner J.S., Ross M. and Hickling N. (1975a) Significance of exsolved pyroxenes from lunar breccia 77215. *Proc. 6th Lunar Sci. Conf.* 529-546.
- Huebner J.S., Ross M. and Hickling N.L. (1975b) Cooling history and significance of exsolved pyroxene in lunar noritic breccia 77215 (abs). *Lunar Sci. VI*, 408-410. Lunar Planetary Institute, Houston.
- LSPET (1973) Apollo 17 lunar samples: Chemical and petrographic description. *Science* **182**, 659-672.
- LSPET (1973) Preliminary Examination of lunar samples. Apollo 17 Preliminary Science Rpt. NASA SP-330. 7-1 – 7-46.
- McGee J.J., Nord G.L. and Wandless M.-V. (1980a) Comparative thermal histories of matrix from Apollo 17 boulder 7 fragment-laden melt rocks: An analytical transmission electron microscopy study. *Proc. 11th Lunar Planet. Sci. Conf.* 611-627.
- McGee J.J., Nord G.L., Jr. and Wandless M.-V. (1980b) Comparative thermal histories of matrix from Apollo 17 boulder 7 fragment-laden melt rocks (abs). *Lunar Planet. Sci. XI*, 700-702. Lunar Planetary Institute, Houston.
- Meyer C. (1994) Catalog of Apollo 17 rocks. Vol. 4 North Massif
- Minkin J.A., Thompson C.L. and Chao E.C.T. (1978) The Apollo 17 Station 7 boulder: Summary of study by the International Consortium. *Proc. 9th Lunar Planet. Sci. Conf.* 877-903.

- Minkin J.A., Thompson C.L. and Chao E.C.T. (1987) Allocation of subsamples of Apollo 17 lunar rocks from the boulder at station 7, for study by the International Consortium. Open-file report 78-511. United States Geological Survey.
- Muehlberger et al. (1973) Documentation and environment of the Apollo 17 samples: A preliminary report. *Astrogeology* 71 322 pp superceded by *Astrogeology* 73 (1975) and by Wolfe et al. (1981)
- Muehlberger W.R. and many others (1973) Preliminary Geological Investigation of the Apollo 17 Landing Site. *In Apollo 17 Preliminary Science Report*. NASA SP-330.
- Nakamura N. and Tatsumoto M. (1977) The history of the Apollo 17 Station 7 boulder. *Proc. 8th Lunar Sci. Conf.* 2301-2314.
- Nakamura N., Tatsumoto M., Nunes P.D., Unruh D.M., Schwab A.P. and Wildeman T.R. (1976) 4.4 b.y.-old clast in Boulder 7, Apollo 17: A comprehensive chronological study by U-Pb, Rb-Sr, and Sm-Nd methods. *Proc. 7th Lunar Sci. Conf.* 2309-2333.
- Nunes P.D., Tatsumoto M. and Unruh D.M. (1974a) U-Th-Pb and Rb-Sr systematics of Apollo 17 Boulder 7 from the North Massif of the Taurus-Littrow valley. *Earth Planet. Sci. Lett.* **23**, 445-452.
- Nunes P.D., Tatsumoto M. and Unruh D.M. (1974b) U-Th-Pb systematics of some Apollo 17 lunar samples and implications for a lunar basin excavation chronology. *Proc. 5th Lunar Sci. Conf.* 1487-1514.
- Nunes P.D., Tasumoto M. and Unruh D.M. (1974c) U-Th-Pb systematics of some Apollo 17 samples (abs). *Lunar Sci.* **V**, 562-564. Lunar Planetary Institute, Houston
- Nunes P.D. and Tatsumoto M. (1975a) U-Th-Pb systematics of selected samples from Apollo 17, Boulder I, Station 2. *The Moon* **14**, 463-471.
- Papike J.J., Fowler G.W. and Schearer C.K. (1994b) Orthopyroxene as a recorder of lunar crust evolution: An ion microprobe investigation of Mg-suite norites. *Am. Mineral.* **79**, 796-800.
- Sanford R.F. and Huebner J.S. (1979) Reexamination of diffusion processes in 77115 and 77215 (abs). *Lunar Planet. Sci.* **X**, 1052-1054. Lunar Planetary Institute, Houston.
- Stettler A., Eberhardt P., Geiss J. and Grogler N. (1974) ³⁹Ar-⁴⁰Ar ages of samples from the Apollo 17 Station 7 boulder and implications for its formation. *Earth Planet. Sci. Lett.* **23**, 453-461.
- Stettler A., Eberhardt P., Geiss J., Grogler N. and Guggisberg S. (1975) Age sequence in the Apollo 17 Station 7 boulder (abs). *Lunar Sci.* **VI**, 771-773. Lunar Planetary Institute, Houston.
- Stettler A. and Albarede Frank (1977) Ar³⁹-Ar⁴⁰ pattern and light noble gas systematics of two mm-sized rock fragments from Mare Crisium (abs). *In Conf. on Luna* **24**. 175-178. Lunar Planetary Institute, Houston.
- Stettler A. and Albarede F. (1978) Ar³⁹-Ar⁴⁰ systematics of two mm-sized rock fragments from Mare Crisium. *Earth Planet. Sci. Lett.* **38**, 401-406.
- Stettler A., Eberhardt P., Geiss J., Grogler N. and Guggisberg S. (1978) Chronology of the Apollo 17 Station 7 Boulder and the South Serenitatis impact (abs). *Lunar Planet. Sci.* **IX**, 1113-1115. Lunar Planetary Institute, Houston.
- Warren P.H. (1993) A concise compilation of petrologic information on possibly pristine nonmare Moon rocks. *Am. Mineral.* **78**, 360-376.
- Winzer S.R., Nava D.F., Schuhmann S., Kouns C.W., Lum R.K.L. and Philpotts J.A. (1974) Major, minor and trace element abundances in samples from the Apollo 17 Station 7 boulder: Implications for the origin of early lunar crustal rocks. *Earth Planet. Sci. Lett.* **23**, 439-444.
- Winzer S.R., Nava D.F., Schuhmann S., Lum R.K.L. and Philpotts J.A. (1975a) Origin of the Station 7 boulder: A note. *Proc. 6th Lunar Sci. Conf.* 707-710.
- Winzer S.R., Nava D.F., Schuhmann P.J., Lum R.K.L., Schuhmann S., Lindstrom M.M., Lindstrom D.J. and Philpotts J.A. (1977b) The Apollo 17 "melt sheet": Chemistry, age and Rb/Sr systematics. *Earth Planet. Sci. Lett.* **33**, 389-400.
- Wolf R., Woodrow A. and Anders E. (1979) Lunar basalts and pristine highland rocks: Comparison of siderophile and volatile elements. *Proc. 10th Lunar Planet. Sci. Conf.* 2107-2130.
- Wolfe E.W., Bailey N.G., Lucchitta B.K., Muehlberger W.R., Scott D.H., Sutton R.L and Wilshire H.G. (1981) The geologic investigation of the Taurus-Littrow Valley: Apollo 17 Landing Site. US Geol. Survey Prof. Paper, 1080, pp. 280.