

Photometric surface properties of Lutetia as seen by OSIRIS/Rosetta. C. Leyrat¹, H. Sierks², A. Barucci¹, S. Fornasier³, M. Fulchignoni³ and the OSIRIS/Rosetta Team, ¹LESIA Observatoire de Paris – Meudon, 5 Place Jules Janssen, F92195 MEUDON France. Email : cedric.leyrat@obspm.fr, ²Max Planck Institut Fur Sonnensystemforschung, 37191 Katlenburg-Lindau, Germany, ³University Paris 7 and Observatoire de Paris.

Introduction: The asteroid (21) Lutetia was successfully observed on July 10, 2010 by the ESA Rosetta spacecraft during its long interplanetary journey toward the comet 67P/Churyumov-Gerasimenko. With a diameter of almost 130 km, Lutetia becomes the largest asteroid observed by a space probe. Because of the trajectory, the northern hemisphere could be observed while the southern remained hidden.

The images obtained in visible wavelengths with the OSIRIS cameras (both NAC and WAC) revealed a variety of craters densities and surface properties, suggesting a different geological history of the areas and different regolith properties. Here we present a spectrophotometric analysis of the surface using the OSIRIS images. Spectro-photometric data were used to generate multi-wavelengths maps of the surface.

Lutetia flyby: During the Lutetia flyby in July 2010, remote sensing instruments onboard the Rosetta spacecraft have acquired precious informations on the surface of the asteroid. The closest approach occurred at 15H45 UTC at a relative speed of 15km/s and a relative distance of 3160 km. The Narrow Angle Camera (NAC) and the Wide Angle Camera (WAC) of the OSIRIS instrument onboard Rosetta acquired images at different phase angles ranging from almost zero to more than 150 degrees. The best spatial resolution (60 m/pixel) allowed to reveal a very complex topography. Numerous sets of images have been obtained at different wavelengths from 270nm to 980nm, allowing to perform a spectro- photometric analysis of the surface. For this purpose, we used 189 images obtained with the NAC with 16 different filters combinations, and 160 images corresponding to 10 WAC combination filters.

Photometric properties: The direct comparison pixel-to-pixel between two images obtained at two different wavelengths and at different time is limited to data acquired at almost the same geometry conditions. At closest approach, a delay of few seconds between two acquisitions imply an important change of spatial resolution and phase angle. Moreover, apparent variation of the surface depends also on shadows and incidence angles that are not representative of the real photometric properties of the regolith.

To generate multi-wavelengths maps of Lutetia, we used the 3D shape model from [1, 2] to create simu-

lated images using the OASIS simulator (LAM). An auto-correlation algorithm was used to adjust OSIRIS data with simulated images on which the local conditions of illumination are known for each pixel (i.e., phase, incidence and emission angle).

Photometric properties were then derive using either [3] photometric law or [4] semi-empirical law. On each case, photometric parameters were first obtained for the full set of pixels on each image by looking for the minimum chi square between the measured and the modeled reflectances. Each image was then corrected assuming a specific phase angle. We will present those images and the areas corresponding to a different photometric properties.

The surface of Lutetia was then analyzed directly facet per facet on the 3d shape model. For each facet located on the northern hemispher, the phase fonction was obtained and photometric parameters were derived independently. We will show the the albedo, roughness and opposition effects parameters and we will discuss the physical implications on the surfaces properties of Lutetia. The northern Beatica region that corresponds to a young surface show important variations of the albedo. We will discuss these variations with the spectral slope. A multivariate statistical analysis (G-mode) will be presented in order to check similar photometric properties accros the surface of the asteroid.

References:

- [1] Jorda L, Lamy P., Groussin O. : 3D shape of Lutetia, in prep.
- [2] Gaskell, R., and 15 co-authors, Characterizing and navigating small bodies with imaging data, Meteoritics & Planetary Science, vol. 43, Issue 6, p.1049-1061, 2008.
- [3] Hapke B, Bidirectional reflectance spectroscopy. 6. Effects of porosity, Icarus, vol 195, p918-926, 2008 2000.
- [4] Akimov L.A., Influence of mesorelief on the brightness distribution over a planetary disk, Sov. Astron. 19, pp. 385–388, 1976