

**Search for Water Vapour Emission from DAWN target (1) Ceres with HERSCHEL.** M. Küppers<sup>1</sup>, L. O'Rourke<sup>1</sup>, D. Bockelée-Morvan<sup>2</sup>, J. Crovisier<sup>2</sup>, B. Carry<sup>1</sup>, D. Teyssier<sup>1</sup>, R. Vavrek<sup>1</sup>, T. G. Müller<sup>3</sup>, M. A. Barucci<sup>2</sup>, B. G. González García<sup>1</sup> and the MACH-11 team

<sup>1</sup>European Space Astronomy Centre, European Space Agency, P.O. Box 78, 28691 Villanueva de la Cañada, Madrid, Spain, [michael.kueppers@sciops.esa.int](mailto:michael.kueppers@sciops.esa.int), <sup>2</sup>Observatoire de Paris, Meudon, France, <sup>3</sup>Max-Planck Institut für Extraterrestrische Physik (MPE), Garching, Germany

**Introduction:** Dwarf planet (1) Ceres is a C-type asteroid [1], associated with the primitive hydrated carbonaceous chondrite CI and CM types. Thermo-physical-chemical models of the evolution of its interior have predicted the possible presence of an ice-rich crust and even liquid water in its interior, with possible polar ice caps [2-5].

Disk-resolved imaging and spectroscopy of Ceres have revealed a surface covered by albedo markings [6-8]. Because water ice is not stable at Ceres' heliocentric distance (2.55 – 2.98 AU), the most sensitive measurement of the presence of (sub-)surface water ice is the search for water vapour or its dissociation products around the asteroid.

In the early 1990s, a marginal (3 sigma) detection of near-ultraviolet OH emission band at 308 nm was indeed reported [9], but could not be confirmed by a later, more sensitive search for the same line [10]. We will present new observations constraining the water production rate of Ceres.

**Observations:** Ceres was observed with the Herodotus Instrument for the Far Infrared (HIFI) on-board Herschel on 23 Nov. 2011 in the context of the Herschel Asteroid and Comet Observation guaranteed time program "MACH 11". The heliocentric and observer distance were 2.94 AU and 2.51 AU, respectively. The water ground state line at 557 GHz was searched for in a 1.1 hour integration. While the thermal continuum from Ceres was detected, no line emission or absorption was found. Additional upper limits of water absorption will be derived from a far infrared spectral range scan of Ceres with the Photodetector Array Camera and Spectrometer (PACS) on Herschel that took place in July 2011 and from further PACS observations taken for Herschel calibration.

**Results and Discussion:** We will model the water line with a "standard" cometary sublimation and radiative transfer model [11], but extending it to include the absorption of the continuum and Ceres' gravity (the escape velocity of Ceres of 0.52 km/s is of the same order as the expansion velocity of cometary water). A preliminary investigation suggests that a water production corresponding to that derived by [9] would have been detected. This implies that the surface fraction covered by water ice is limited to a maximum of order

$10^{-7}$ , consistent with the non-detection by [6-8]. If Ceres crust indeed contains large amounts of water ice, it must be covered by a thick dust layer.

A deeper search for water vapour on Ceres will be performed by Herschel in winter 2012/2013, closer to perihelion. Notably, both the measurements in [10] and the present observations were performed at larger heliocentric distance than the tentative detection by [9]. If the different measurements are indicative of a strongly varying activity of Ceres, there is a good chance of a detection in the upcoming observation. The question of the presence of water ice on Ceres may be finally answered in 2015 when it will be encountered by the DAWN spacecraft.

#### References:

- [1] DeMeo et al., 2009, *Icarus* 202, 160-180.
- [2] Fanale & Salvail 1989, *Icarus* 82, 97-110.
- [3] McCord & Sotin, 2005, *JGR* 110, 5009-5023.
- [4] Zolotov, 2009, *Icarus* 204, 183-193.
- [5] Castillo-Rogez & McCord, 2010, *Icarus* 205, 443-459.
- [6] Li et al., 2006, *Icarus* 182, 143-160
- [7] Carry et al., 2010, *A&A* 478, 235-244
- [8] Carry et al., 2012, *Icarus* 217, 20-26
- [9] A'Hearn, & Feldman 1992, *Icarus* 98, 54-60.
- [10] Rousselot et al. 2011, *Astron. J.* 142, 125.
- [11] Bockelée-Morvan & Crovisier 1987, *A&A* 181, 169-181.