

LABORATORY SIMULATIONS OF LARGE-SCALE VORTEX FLOWS GENERATED AT IMPACTS ON VENUS AND ON EARTH. V.I.Artem'ev, V.A.Rybakov, S.A.Medveduk, and B.A.Ivanov. Institute for Dynamics of Geospheres, Russian Academy of Sciences, Leninsky prosp., 38-6, Moscow, 117979.

The analysis of the Magellan Mission data suggests that ejecta emplacement and crater formation on Venus are closely related to the atmospheric effects. The formation of the parabolic impact crater related features is supposed to be determined to the injection of small particles to the upper atmosphere and their transport by E-W zonal winds. We propose here another model of ejecta long distant transportation: the escape of ejecta particles to high altitudes can be performed by large-scale atmospheric vortex flows.

Collisions of large meteoroids with planets having dense atmospheres (Venus and Earth) are accompanying by substantial atmospheric disturbances. The impact-generated atmospheric flows with vertical size exceeding the scale height of the atmosphere may be responsible for global propagation of solid ejecta and dust.

In [1] a model for the formation of the parabolic impact crater related features is developed based on the injection of small particles to the upper atmosphere. The vorticity generation at impacts on Venus has been attributed to the interaction of atmospheric blast waves with radiatively heated surface (the thermal layer effect) [2], and to the atmospheric disturbances by the outward moving ejecta [3]. In [4,5] it was emphasized that a disturbance of the atmosphere by falling meteoroid *before* the impact (i.e. a formation of the wake of shock-heated gas downstream of the meteoroid) promotes the escape of the ejecta to the upper layers of the Venusian atmosphere; in experiments [6] the near-surface vortex flow has been generated by detonation of an oblique line charge burst simulated the atmospheric wake.

In general, the baroclinity of the gasdynamic flow ($\nabla 1/\rho \times \nabla p \neq 0$, where ρ and p are density and pressure) is responsible for vorticity generation. A complex time-spatial character of energy release during atmospheric flight and at impact of the meteoroid - the inhomogeneity of the atmosphere, the presence of heated channels and layers, crossing shock fronts *etc.* - results in such a baroclinity and hence in vorticity generation. In our experiments we make an effort to simulate the vortex flows resulting from the interactions of impact-generated air blast wave and wake.

The spherical shock wave is produced on action of a focused laser radiation onto a surface. The energy of the laser driven blast is $\mathcal{E} \approx 10$ J. The atmospheric wake is simulated by electrical explosion of a thin wire. On modeling of the impact on the Earth the atmospheric pressure and density in laboratory are the same as in real atmosphere. In this case the energy \mathcal{E} corresponds to the energy of impact-generated air blast E as follows: $\mathcal{E} = (l/L)^3 \cdot E$, where l and L are the linear scales in laboratory and in atmosphere. The explosion energy for 100 m-stony asteroid falling with velocity ~ 15 km/s is about $E \approx 10^{24}$ erg. In this case $L = 10$ km in atmosphere corresponds to $l \approx 5$ cm in laboratory. The energy expended for formation of the atmospheric wake can be estimated in the case considered as $E_a \approx 2\rho_a H / (\rho_m D) \cdot E \approx 0.1 \cdot E$, where $H \approx L = 10$ km is the scale height of the Earth's atmosphere, and $D = 100$ m is the asteroid's diameter. Thus the energy of electrical explosion simulating the wake must be $\mathcal{E}_a / l \approx 0.1 \cdot \mathcal{E} / l = 0.2$ J/cm.

Fig.1 represents the results of the laboratory simulation of the impact on the Earth: the shadowgraphs of the flow generated on blast-wake interaction at the consecutive points of time. The "experimental" time τ corresponds to the "real" time t as $\tau = l/L \cdot t \approx 5 \cdot 10^{-6} \cdot t$. In Fig.1a one can see the expanding wake just before the impact, two next pictures illustrate the

LARGE-SCALE VORTEXES AT IMPACTS: V.I.Artem'ev *et al.*

interaction of spherical and cylindrical air blast waves. In Fig.1d and Fig.1e are shown the initial stages of the large-scale vorticity formation: initially the near-surface part of the wake and then the whole wake is destroyed by vortexes. On the later stage (Fig.1f, $\tau=5$ ms, $t=17$ min) a system of large-scale toroidal vortexes oriented normally to the initial direction of the wake is formed.

In simulation of the impact of 3 km-projectile on Venus the energy of atmospheric explosion is taken to be comparable to the energy dissipated in the wake. In this case the results are qualitatively the same as for the impact on the Earth. Notice that in both cases we neglect the atmospheric inhomogeneity. The vertical scale of vortex flows (especially at impact on Venus) in fact exceeds the scale height of atmospheres, therefore in more improved simulation a vertical decrease of the atmospheric density will be taken into account.

References: [1] Campbell D.B. *et al.* (1992), J. Geophys. Res., 97, No E10, p.16249-16277. [2] Ivanov B.A. *et al.* (1992), LPSC XXIII, p.575-576. [3] Barnouin O.S. and Schultz P.H. (1993), LPSC XXIV. [4] Ivanov B.A. *et al.* (1986), J. Geophys. Res., 91, No B4, p.413-430. [5] Ivanov B.A. *et al.* (1992), J. Geophys. Res., 97, No E10, p.16167-16181. [6] Ivanov B.A. and Provalov A.A. (1993), LPSC XXIV.

Fig.1

