

EXPERIMENTAL DETERMINATION OF SPINEL/MELT, OLIVINE/MELT, AND PYROXENE/MELT PARTITION COEFFICIENTS FOR Re, Ru, Pd, Au, AND Pt. K. Righter¹, A.J. Campbell² and M. Humayun²; ¹NASA Johnson Space Center, Mail Code ST, Houston, TX 77058, kevin.righter-1@nasa.gov; ²Department of the Geophysical Sciences, The University of Chicago, 5734 S. Ellis Ave., Chicago, IL 60637.

Introduction

Experimental studies have identified spinels as important host phases for many of the highly siderophile elements (HSE) [1-4]. Yet experimental studies involving chromite or Cr-rich spinel have been lacking. Experimental studies of partitioning of HSEs between silicate, oxides and silicate melt are plagued by low solubilities and the presence of small metallic nuggets at oxygen fugacities relevant to magmas, which interfere with analysis of the phases of interest [5]. We have circumvented these problems in two ways: 1) performing experiments at oxidized conditions, which are still relevant to natural systems but in which nuggets are not observed, and 2) analysis of run products with laser ablation inductively coupled plasma mass spectrometry (LA-ICP-MS), allowing a combination of high sensitivity and good spatial resolution.

Experimental and analytical

Partition coefficients for Re, Ru, Pd, Au and Pt have been measured using three different compositions. Ankaramite, doped with Cr₂O₃, encapsulated with AuPd tubing, and buffered with Fe₂O₃-Fe₃O₄ and MnO-Mn₃O₄, was used to study spinel/melt and olivine/melt partitioning. Two synthetic basalts, encapsulated by Pt and buffered with Ni-NiO and Ru-RuO₂, were used to study spinel/melt and clinopyroxene/melt partitioning. The 1 bar experiments were carried out in evacuated silica tubes, and the 20 kbar experiments were carried out in a piston cylinder apparatus. An electron microprobe was used to analyze major elements in all phases, and Ru and Rh in some spinels. Highly siderophile elements were measured by LA-ICP-MS, using

100 μm, 50 μm and 25 μm spot sizes, respectively, in glass, olivine and spinel. HSE abundances were determined relative to MgO contents of the phases, which were analyzed independently by electron microprobe. The absence of nuggets was inferred from the smooth quality of the time dependent LA-ICP-MS signals, the reproducibility of analyses in adjacent regions of the glass, and the agreement with HSE solubility studies [6-11].

Results

When our new results for Ru (Table 1) are combined with previous results [1], it is evident that spinel/melt partition coefficients for Ru are highest at the lowest oxygen fugacities ($D = 1100-2300$), and/or when the spinel is inverse such as magnetite (# 118; $D = 870$). These results suggest that both inverse and chromian spinels can be important hosts for Ru at mantle oxygen fugacities, in agreement with partition coefficients inferred from komatiite lava lakes ($D(\text{Ru})$ spinel/melt = 151; [12]). On the other hand, Re is incompatible in all spinels produced experimentally so far. These trends may reflect higher valences for Re and Ru at oxidized conditions (6+ and 4+), compared to reduced conditions (3+?). Platinum is compatible, whereas Re is incompatible in clinopyroxene (Table 1). Ruthenium, Au and Pd are incompatible in olivine (Table 1), in agreement with previous experimental work [1,3].

Applications

Correlations of Ru, Ir and Rh with Cr and MgO observed in komatiites and basaltic komatiites from Kostomuksha (Figure 1) [12], Cape Smith [13], and the Vetryny Belt [14] can all be explained by a combination of oli-

Spinel/melt partitioning of HSE. K. Righter et al.

vine and chromite fractionation, using the partitioning results from this study. Incompatibility of Re in oceanic basalt suites [15,16] is also supported by our measured $D(\text{Re})$ spinel/melt $\ll 1$. In addition, the mildly compatible nature of Pt in clinopyroxene ($D(\text{Pt})$ cpx/melt = 1.5), together with the affinity of Pt for sulfide melt [17,18] can explain the very similar Pt concentrations observed in martian clinopyroxeneites (sulfide-bearing; [19]) and shergottite basalts. Application of these new results will also be made to Apollo 12 and 15 basalt suites.

This research is supported by NASA grant NAG5-9435 and NSF grant EAR0074036.

Table 1: Summary of experimental conditions and partition coefficients

run	158	118	153
phase	sp, ol	sp	cpx
P, T (°C)	1 bar, 1300	20 kb, 1275	1 bar, 1300
buffer	Fe ₂ O ₃ -Fe ₃ O ₄	Ru-RuO ₂	Ni-NiO
logfO ₂	-1.5	-1.3	-6.7
D(Re)	-	-	0.18
D(Ru)	76, 0.24	870	-
D(Pd)	0.14, 0.12	-	-
D(Pt)	-	-	1.50
D(Au)	0.08, 0.13	-	-

References:

[1] Capobianco, C. J., R. L. Hervig and M.J. Drake (1994) *Chemical Geology* 113, 23-43; [2] Capobianco, C. J. and M. J. Drake (1990) *Geochimica et Cosmochimica Acta* 54, 869-874; [3] Capobianco, C. J., M.J. Drake, and Rogers, P.S.Z (1991) *Lunar Planet. Sci. XXII*, 179-180; [4] Righter, K. and R.T. Downs (2001) *Geophysical Research Letters* 28, 619-622; [5] Lindstrom, D.J. and Jones, J.H. (1996) *Geochimica et Cosmochimica Acta* 60, 1195-1203; [6] Borisov A. and Palme H. (1995) *Geochim. Cosmochim. Acta* 59, 481-485; [7] Borisov, A. and H. Palme (1996) *Mineralogy and Petrology* 56, 297-312; [8] Borisov, A. and Nachtweyh, K. (1998) *Lunar. Planet. Sci. XXIX*, abstract #1320; [9] Ertel, W., O'Neill, H.St.C., Sylvester, P.J. and Dingwell, D.B. (1999) *Geochimica et Cosmochimica Acta* 63, 2439-2449; [10] Ertel, W., O'Neill, H.St.C., Sylvester, P.J., Dingwell, D.B. and B. Spettel (2001) *Geochimica et Cosmochimica Acta* 65, 2161-2170; [11] Righter K, Drake MJ. (1997) *Earth Planet.*

Sci. Lett. 146:541-53; [12] Puchtel I. and Humayun, M. (2000) *Geochimica et Cosmochimica Acta* 64, 4227-4242; [13] Barnes, S.-J. and Picard, C.P., 1993. *Geochim. Cosmochim. Acta* 57, 79-87; [14] Puchtel I. and Humayun, M. (2001) *Geochimica et Cosmochimica Acta* 65, 2979-2993; [15] Hertogen J., Janssens M.-J. and Palme H. (1980) *Geochim. Cosmochim. Acta* 44, 2125-2143; [16] Righter, K., Chesley, J.T., Geist, D. and Ruiz, J. (1998) *Journal of Petrology* 39, 785-795; [17] Fleet, M. E., W. E. Stone, et al. (1991) *Geochimica et Cosmochimica Acta* 55, 2545-2554; [18] Fleet, M. E., J. H. Crocket, et al. (1996). *Geochimica et Cosmochimica Acta* 60, 2397-2412; [19] Bunch, T.E., and A.M. Reid (1975) *Meteoritics* 10, 303-315; [20] Ghiorso M. S. and Sack R. O. (1994) *Contrib. Mineral. Petrol.* 119, 197-212.




Figure 1: Cr, Ru and Pd trends with MgO in the Kostomuksha komatiite suite (from [12]). Major element modelling of crystal fractionation of the proposed Kostomuksha parent liquid [12] using the MELTS program [20] is coupled with new partitioning data. Asterisks indicate the calculated trends using the major element results, and $D(\text{Ru})$ and $D(\text{Pd})$ olivine/melt and chromite/melt measured here.