

THERMOPHYSICAL, RHEOLOGICAL AND MECHANICAL MEASUREMENTS ON ICY COMPOSITIONS WITH APPLICATION TO SOLAR SYSTEM ICES. C. C. Hays¹, J. C. Castillo¹, K. L. Mitchell¹, M. Barmatz¹, F. Zhong¹, W. Smythe¹, D. L. Matson¹, R. T. Pappalardo¹, R. M. C. Lopes¹, L. E. Robshaw^{3,1}, C. Neish⁴, J. I. Lunine⁴, and J. S. Kargel², ¹Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Drive, Pasadena, CA 91109. E-mail: Charles.C.Hays@jpl.nasa.gov, ²Department of Hydrology and Water Resources, Univ. Arizona, ³Lancaster University, Environmental Sci. Dept., Lancaster, UK, ⁴Dept. of Planetary Sci., University of Arizona.

Introduction: It is necessary to better understand the properties of the icy materials that make up ice-rock planetary bodies in order to fully appreciate the discoveries made by the *Cassini-Huygens* and *Galileo* missions, to prepare for the *Dawn* and *New Horizon* mission, and to look forward to potential missions to Europa, Enceladus, and Titan. We present the motivations and objectives of a new experimental cryo-ices initiative at JPL. This is a joint effort among experimentalists and theorists at JPL, in collaboration with specialists in ice properties.

Science Motivation: Differences in geomorphology among the surfaces of the Galilean and Saturnian satellites on one hand and the Uranian and Neptunian satellites on the other hand have been attributed to differences in ice composition [1]. To a large extent this is true for differences among morphologies within these systems as well.

A range of theoretical needs and observations have led us to develop an experimental initiative aimed at understanding the link between ice composition (including structure) and icy satellite observations.

Icy Satellite Internal Evolution. Internal evolution models depend upon accurate characterization of the elastic moduli, deformation pathways and mechanisms, and solid-state rheology of the component materials. Ice rigidity and viscosity play a role in the modeling of convection, differentiation, geological activity (lithospheric response to loading, crater relaxation, folding/faulting as a function of strain rate), ice behavior at premelting conditions, and cryovolcanism. The role of ice properties in many of these processes is addressed by [2]. Also these properties play a major role in the tidal response of the bodies, and associated tidal dissipation. For the icy moons of the outer satellites, cosmochemical arguments suggest that these materials will be dominated by water, with variable amounts of materials plausibly including methane clathrate hydrates, ammonia, methanol [3, 4], and salts [5].

Enceladus' Resurfacing Processes. A major discovery of the Cassini-Huygens mission is the observation of a hot region surrounding Enceladus' South pole, associated with a surface likely less than 1 My old, faults (the "tiger stripes"), and intense plume-generating activity [6,7]. Plausibly cryovolcanic landforms have been identified on Titan [8-10], which may

be ammonia hydrate slurries and which are associated with methane, and possibly containing organics. Cryovolcanic activity has been suggested to be responsible for resurfacing and/or landform construction on the surfaces of Ganymede, Miranda, Ariel, Triton, and others, driven by radiogenic heating and/or tidal heating [11]. However, our understanding of mechanisms behind these processes is limited; in particular, how ice responds mechanically to tidal stresses, and the behaviour of eruptive materials during flow, are poorly constrained.

Titan's Complex Environment. Titan's atmospheric and crustal volatile chemistry produce a complex assemblage of materials, in some cases beyond those of other icy bodies, that may be mobilized superficially and in the upper crust [12], and could exhibit a wide range of behaviours. These may play an important role in limnologic (lake) [13], fluvial, sedimentary, aeolian [14], glacial and periglacial processes [12,15]. Despite some successful characterization of Titan's surface and atmospheric materials by Cassini VIMS spectral analysis [7], it seems unlikely that Cassini will be able to determine much of the range of important surface materials through Titan's obfuscating atmosphere. Therefore, in order to understand the materials involved in the various processes on the surface of Titan, we need to use a less-direct approach.

We use combinations of observation, theory and experiment to relate primarily morphological observations of surface features to the processes and materials that could have formed them. On Titan, such materials most likely include liquid methane and/or ethane, cryovolcanic ammonia hydrate slurries (possibly containing organics), and a variety of carbonaceous compounds, organics, tholins, salts and hydrates.

In the above cases, our ability to relate landforms and other observations to the processes that formed them is sorely limited by physical models that contain incomplete databases of the physical properties of the candidate materials. These properties include the rheological behaviour of candidate cryovolcanic slurries, the loss tangent of liquid methane/ethane, rates of dissolution and wetting angles of liquid methane and ethane on candidate surface materials, and the mechanical properties and mobility of surface solids (water-ice, ammonia hydrates, acetylene, etc.).

Approach: A range of experiments is being devised which will improve our ability to model ice-rock body internal evolution and geological processes using modern synthesis and characterization techniques under cryogenic conditions. Initial experiments will involve pure water (H₂O), methanol-water (CH₃OH-H₂O), ammonia-water (NH₃-H₂O) and ammonia-water-methanol mixtures, relevant to a range of icy satellites and processes. Where beneficial, we will determine basic thermophysical properties and phase diagrams using a Differential Scanning Calorimeter.

Solids. Our specimen synthesis conditions for solids are designed to mimic processes that may be part of the formation of icy planetary landscapes. The synthesis methods for icy specimens will exploit equilibrium and non-equilibrium methods. Equilibrium methods, *e.g.*, conventional (slow) freezing in a mold, will provide microstructural length scales, *d*, in the range ~200 μm to 10 mm (with and without preferred orientation, *e.g.*, columnar grains). Non-equilibrium methods, *e.g.*, rapid-quenching, will vitrify the specimens, and we will control the specimen microstructure (*e.g.*, grain size) via low temperature annealing.

Post-synthesis microstructural characterization will be performed using Cryogenic Optical Microscopy integrating a cross-polarizer to analyse thin sections, and a Cryogenic Scanning Electron Microscope.

Mechanical properties of solid specimens will be performed for temperatures between 80 and 270 K with a cryogenically cooled Instron measurement system. Compression measurements will be conducted as a function of temperature, strain-rate, microstructural length scale and orientation, applied load versus time (creep), time-varying load in order to measure the viscous response. The cryogenic stage for use with the Instron test platform is in development. We have begun testing and calibrating the Instron using Teflon and ice samples.

Terrestrial glaciers appear as realistic analogs for modeling processes taking place in icy satellite outer icy shells [6,19-21]. As a follow-on to our initial studies, we plan to analyze samples from terrestrial glaciers, which are relevant to both the terrestrial and planetary geology and geophysical communities. These results will better enable us to predict the long-term evolution of terrestrial glaciers and ice shelves, and to provide support material to infrared observations.

Fluids. In order to improve our understanding of effusive cryovolcanism, we will measure rheological properties of liquid and mixed (slurry) materials between 80 and 300 K using a cryogenically cooled Brookfield rotational rheometer, based on the experi-

mental procedures of Kargel *et al.* [22], but over a much wider range of parameters and with a greater degree of automation. Control and data acquisition will be carried out using a LabView custom interface. The results will be integrated into parallel modeling efforts that include the slurry rheology modeling discussed above.

Methane wetting and dissolution of water-ice and other materials will be carried out in order to explore the effects of the presence of methane lakes on Titan's surface.

Follow-on experiments will explore the dissipation factor at low frequencies approaching satellite orbital frequencies. We will investigate more complex materials relevant to surface processes on Titan, including hydrocarbons such as acetylene and benzene, as well as tholins and clathrates, which should exhibit a range of rheological and mechanical properties from fast-moving fluids to glacial creep.

Acknowledgements: Much of the research described in this presentation was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration. KLM is supported by a NASA Postdoctoral Fellowship, administered by Oak Ridge Associated Universities. JCC is a postdoctoral researcher of the California Institute of Technology.

References: [1] Kargel J. S. (1994) *Earth Moon & Planets*, 67, 101-113. [2] Durham and Stern (2001) *Ann. Rev. Earth Planet. Sci.*, 29, 295-330. [3] Tobie G. *et al.* (2005) *Icarus*, 175, 496-502. [4] Prieto-Ballesteros *et al.* (2005) *Icarus*, 177, 491-505. [5] Kargel J. S. (1991) *Icarus*, 94, 368-390. [6] Spencer *et al.* (2006) *Science*, 311, 1401-1405. [7] Porco *et al.* (2006) *Science*, 311, 1393-1401. [8] Elachi C. *et al.* (2005) *Science*, 308, 970-974. [9] Sotin C. *et al.* (2005) *Nature*, 435, 786-789. [10] Lopes R. M. C. *et al.* (2007) *Icarus*, in press. [11] Schenk, P. M. and Moore, J. M. (1998). *Solar System Ices*, 551-578. [12] Kargel J. S. *et al.* (2007) *LPSC XXXVIII*, Abstract, this volume. [13] Stofan E. R. *et al.* (2007) *Nature*, 445, 61-64. [14] Lorenz R. D. *et al.* (2006) *Science*, 312, 724-727. [15] Mitchell K. L. *et al.* (2007b) *LPS XXXVIII*, Abstract, this volume. [16] Clark R. N. *et al.* (2006) *Eos. Trans. AGU 87(52)*, Fall Meet. Suppl., Abstract P11A-03. [17] Sandwell S. *et al.* (2004) *JGR*, 109, E11003. [18] Duval P. and Montagnat M. (2004) *Workshop on Europa's Icy Shell*, Abstract #7001. [19] Barr A. and McKinnon W. B. (2007) *JGR* in press. [20] Kargel *et al.* (1991) *Icarus*, 89, 93-112.