

STRATIGRAPHY AND STRUCTURE OF THE NORTH POLAR LAYERED DEPOSITS ON MARS. K. E. Herkenhoff, C. Fortezzo, G. Cushing, R. L. Kirk, L. A. Soderblom, and L. Weller, U. S. Geological Survey Astrogeology Science Center, 2255 N. Gemini Dr., Flagstaff, AZ 86001 (kherkenhoff@usgs.gov).

Introduction: The Martian polar layered deposits (PLD) have long been thought to be the best source of information about the recent climate history of Mars [1-3], but the detailed mechanisms of accumulation are still a mystery [4]. The polar layers are sedimentary deposits composed of water ice and varying amounts of dust or other impurities [5]. Because climate changes are likely recorded as variations in composition or deposition/erosion rates between layers, the detailed stratigraphy of the PLD is of great interest. Layer thicknesses of ~10 to 50 m were observed in Viking Orbiter images of the north PLD [6], and Mars Orbiter Camera (MOC) images resolve layers with similar or lesser thicknesses in both polar regions [7]. In order to accurately determine the thickness of layers and interpret PLD stratigraphy and structure, the topography of exposures must be known. Previous studies have modeled brightness variations in the PLD in an attempt to link them to orbital/axial variations [8-10], but lack of detailed topography has hindered stratigraphic interpretations. More recently, HiRISE images have shown that brightness variations between layers can be caused by mantling deposits of frost and other (dusty) material and are therefore probably not related to the internal composition of the layers [11]. Therefore, brightness variations cannot be used alone to infer PLD stratigraphy; high-precision topographic information is needed. Here we describe results of our continuing study to evaluate the topography, structure, and stratigraphy of the north PLD using photogrammetry on MOC images.

Approach: We used a 2-dimensional photogrammetric technique [12] constrained using simultaneously-acquired MOLA data. This technique is well suited to images taken at high latitudes when the surface was covered by seasonal frost and the solar elevation angle was low so that albedo variations and their effects are minimized and topographic modulation is emphasized. The high density of MOLA data in the polar regions allows gridded topographic products to be generated at higher spatial resolution (~250 m/pixel) than is possible at lower latitudes (Fig. 1). The MOLA gridded data are used to initialize the photogrammetric solution and constrain surface and atmospheric scattering parameters. The ability to discern the layers facing the sun in the photogrammetrically-derived digital elevation model (DEM) is at essentially the same spatial resolution as the original image.

Results: Figure 2 shows some topographic profiles across the layers shown in Figure 1. Individual layers

visible in the MOC images are typically 20-40 m thick, as noted in previous studies. Many layers are characterized by a raised “rim” or “lip.” While it is possible to correlate stratigraphy in the data shown here, correlation is difficult in other areas due to unconformities and other structural complexity. Further analysis of MOC photogrammetric models of this and other PLD exposures will be presented at the conference.




Figure 1. MOC images on MOLA shaded relief, with illumination from left. Yellow dots represent starting locations of layer measurements shown in top part of Fig. 2. Red line shows location of cross-section displayed in bottom part of Fig. 2.

References:

- [1] Murray, B. C., *et al.* (1972). *Icarus* **17**, 328-345.
- [2] Cutts, J. A., *et al.* (1979). *J. Geophys. Res.* **84**, 2975-2994.
- [3] Thomas, P., *et al.* (1992). In *Mars*, University of Arizona Press, Tucson, pp. 767-795.
- [4] Byrne, S. (2009). *Ann. Rev. Earth Planet. Sci.* **37**, 535-560.
- [5] Phillips, R. J., *et al.* (2008). *Science* **320**, 1182-1185.
- [6] Blasius, K. R., *et al.* (1982). *Icarus* **50**, 140-160.
- [7] M. C. Malin and Edgett, K. S. (2001). *J. Geophys. Res.* **106**, 23,429-23,570.
- [8] Laskar, J., *et al.* (2002). *Nature* **419**, 375-377.
- [9] Milkovich, S. M. and J. W. Head III (2005). *J. Geophys. Res.* **110**, 1005.
- [10] Perron, J. T. and P. Huybers (2008). *Geology* **37**, 155-158; doi: 10.1130/G25143A.1.

[11] Herkenhoff, K. E., *et al.* (2007). *Science* **317**, 1711-1715.

[12] Kirk, R. L., *et al.* (2003). ISPRS Working Group IV/9 Workshop "Advances in Planetary Mapping 2003", Houston.




Figure 2. (top) Layers have been aligned in the location of the most reasonable correlation (boxed area). Other correlations are made based on similar stratigraphic patterns. This method does not account for lateral thickness variations and ignores thinner units. (bottom) Fence diagram showing layers in alternating colors. Light and dark blue layers are those correlated in the plot above. Dashed lines represent likely location of layers in the intermediate trough, interpolated linearly. The colors of the MOLA topographic profile represent the ABb1 (red) and ABb2 (blue) units from Tanaka *et al.*, 2008. Profile is ~23 km long, vertical exaggeration ~5X.