

LUNAR ORBIT ANOMALY AND $GM=tc^3$ COSMOLOGY. L. M. Riofrio¹, ¹University of Houston Clear Lake, Sailorstarfightr@yahoo.com

Introduction: Studies of the Moon, made thanks to Johnson Space Center, have confirmed a large anomaly in lunar orbital evolution. The Lunar Laser Ranging Experiment (LLRE) has reported the Moon's semimajor axis increasing at 3.82 ± 0.07 cm/yr, anomalously high. Tidal data indicates a recession rate of only 2.9 ± 0.6 cm/yr. Additional observations independently measure a recession rate of 2.82 ± 0.08 cm/yr. A cosmology where speed of light c is related to time t by $GM = tc^3$ has been suggested to predict the redshifts of Type Ia supernovae. By this hypothesis, lunar orbital distance would appear to increase an additional 0.935 cm/yr. An anomaly in the lunar orbit may be precisely accounted for, shedding light on puzzles of "dark energy." This hypothesis may also explain the "faint young sun" paradox of astrophysics.

Lunar Orbit Anomaly: The Moon has long been known to be slowly drifting from Earth due to tidal forces. As the Moon raises tides on Earth, tidal bulges travel ahead of the Moon due to Earth's 24-hour rotation. A tidal bulge pulls the Moon slightly ahead in its orbit, and causes Earth's rotation rate to slow. In this way angular momentum is transferred from Earth's rotation to the Moon, causing the Moon's semimajor axis to increase.

The Lunar Laser Ranging Experiment bounces light off corner reflectors left on the Moon by Apollo astronauts. Previously LLRE has been used to investigate geophysics of the Earth-Moon system and test Relativity. Accuracy is considered fine enough to constrain changes in Newton's gravitational constant G . LLRE has measured the Moon's semimajor axis at $a = 384,402$ km. Repeated measurements appear to indicate that distance increasing 3.82 ± 0.7 cm/yr, anomalously high. [1] If the Moon were gaining angular momentum at this rate, it would have coincided with Earth less than 2 Gyr ago. Our studies of lunar samples convincingly show that the Moon has existed separately from Earth over 4.5 Gyr.

Tidal Rhythmites: Geology and paleontology can also tell how the Moon's distance has changed. Tidal rhythmites, in particular, carry a record of lunar-induced tides. Thicknesses of sedimentary layers vary with the height of local tides. Rhythmites can be used to determine lunar distance over hundreds of millions of years.

Starting with today's LLRE measurement, we may compile estimates of lunar orbital distance based on rhythmites. [2]

Sediment Location	Age 10^3 yr	Distance 10^3 km
Present	0	384.4
Mansfield	310 ± 5	375.3 ± 1.9
Elatina	620 ± 100	370.9 ± 0.1

The Mansfield sediment of Indiana, the most recent, places the Moon $375,300 \pm 1,900$ km away 310 Myr ago, a recession rate of 2.9 ± 0.6 cm/yr. An independent study of the Elatina and Reynella tidal rhythmites also indicates a lower recession rate. [3]

Eclipse Records: Corroborating data may have come from historical astronomers. If the narrow track of total eclipse is recorded over an observatory, it provides an accurate measure of Earth's slowing rotation rate. As Earth and Moon form a closed system, this tells us how much angular momentum has been transferred. A recession rate of 3.82 ± 0.7 cm/yr corresponds to change in Earth's length of day of 2.30 msec/cyr. Observations spanning 2700 yr show change in LOD of $1.70 \pm .05$ msec/cyr, corresponding to a lunar recession of 2.82 ± 0.08 cm/yr. [4] Though eclipse records corroborate tidal data, LLRE's laser light differs by over 10σ .

Speed of Light: As with Mercury, anomalies in orbits may have implications for Physics. One theory of Space/Time suggests that speed of light c is related by:

$$GM = tc^3$$

Where t is age of the Universe, GM combines its mass and gravitational constant. [5]

Solving, we would have:

$$c(t) = (GM)^{1/3} t^{-1/3}$$

Time for light to return would increase yearly, making the Moon appear to recede faster as measured by LLRE.

This model has previously been suggested to predict "accelerating" redshifts of Type Ia supernovae. It may also be used to model features seen in the cosmic microwave background, predicting a 4.507034% proportion of baryonic matter.

Apparent anomaly in lunar distance would then be proportional to change in c :

$$\frac{\dot{a}}{a} = -\frac{\dot{c}}{c} = \frac{1}{3t}$$

Multiplied by the Moon's semimajor axis, that distance would appear to increase an additional 0.935 cm/yr. Theory precisely accounts for the 10σ anomaly.

Faint Young Sun: The speed of light may help explain a puzzle of solar luminosity. Astrophysicists' models predict that, other factors being equal, the early Sun would have only 75% of its present luminosity. Earth's average temperature would have been only 263K, below freezing point. Appearance of life and its evolution would have been unlikely.

Geology shows evidence of sediments and liquid water on Earth at 4 Gyr. Paleontology dates the earliest organisms at least 3.4 Gyr. Spacecraft images show signs of flowing water on Mars at 3.8 Gyr. Evidence from Martian meteorites also shows conditions suitable for liquid water and life. Conflict of astrophysical hypothesis with reality is called the "faint young sun" paradox. [6]

A solution may be in the speed of light. The Sun converts fuel to energy according to $E = mc^2$. Earth is estimated to be 4.6 Gyr and the Universe 13.7 Gyr old, 1.5 times its age at Earth formation. Billions of years ago, solar luminosity may have been higher than once thought.

Solar output would then be adjusted by:

$$\left(\frac{c_i}{c_0}\right)^2 = \left(\frac{t_0}{t_i}\right)^{2/3} = 1.31$$

Multiplied by an initial estimate of 75%, luminosity at Earth formation was 98% of today's value.

Extrapolating through Earth history, the "solar constant" indeed appears nearly constant. The narrow range of temperatures necessary for liquid water constrains solar luminosity. If c had not changed in the amounts predicted, life might not have evolved on Earth.




Fig. 1: Solar luminosity vs. solar system age. L/L_0 is luminosity as a fraction of present value. Lower line is standard solar model. Upper line indicates luminosity when c change is a factor. If speed of light c is precisely related to Universe age t by $GM = tc^3$, luminosity remains nearly constant.

Discussion: At one time controversy raged whether light was instantaneous or had a finite speed. Galileo reportedly suggested measuring the velocity of light with lanterns placed on hilltops. Today we have laser lanterns and the distant hill of the Moon. The Lunar Laser Ranging Experiment, considered highly accurate, may reveal possible variation in c .

The 2.9 ± 0.6 cm/yr recession rate found in Mansfield sedimentary data has been attributed to anomalous tides over many millions of years, an inference that is not independently verified. The lower recession rate is verified by the more precise 2.82 ± 0.08 cm/yr found from eclipse data. An anomaly in eclipse records just 2700 yr old is convincingly non-tidal in origin. Corrected for the speed of light, lunar recession rate would be 2.88 ± 0.07 cm/yr. This value is in agreement with sedimentary data, eclipse records and the Moon's geologic age.

A predicted change in the speed of light may have been seen in redshifts of Type Ia supernovae. Rather than acceleration, redshifts may show signs of c slowing. Supernova observations may closely corroborate lunar data.

The speed of light has been a subject of speculation since at least the time of Tait and Thomson [7]. More recently varying c has been promoted by Moffatt [8], Albrecht and Maguiejo [9]. Experiments involving the Sun, Moon and supernovae, when viewed together, may indicate a "c change" in physics. In Planck units $GM = tc^3$ may be expressed as $M = R = t$.

This work was assisted by NASA and serves goals of the Beyond Einstein program. Great thanks are due to colleagues at the ARES group in Johnson Space Center, Houston. Valuable discussions were made with Marni Dee Sheppard and Yves-Henry Sanejouand.

References:

- [1] Dickey J. O. et al. (1994) *Science*, 265, 482.
- [2] Bills A. B. and Ray C. D. (1999) *Geophysical Research Letters*, 26, 3045.
- [3] Williams G. E. (2000) *Reviews of Geophysics*, 38, 37.
- [4] Stephenson F. R. and Morrison L. V. (1995) *Phil. Trans. R. Soc.*, 351, 165.
- [5] Riofrio L. M. (2005) *Observing Dark Energy*, 339, 181.
- [6] Sagan C. E. and Mullen G. (1972) *Science*, 177, 52.
- [7] Thomson W. and Tait P. G. (1874) *Natural Philosophy*, 1, 403.
- [8] Moffat J. and Clayton M. (1999) *Physics Letters B*, 460, 263.
- [9] Albrecht A. and Maguiejo J. (2000) *Phys Rev D*, 63, 52.