


Measurements of Mars Methane at Gale Crater by the SAM Tunable Laser Spectrometer on the Curiosity Rover

Chris R. Webster¹, Paul R. Mahaffy², Sushil K. Atreya³, Greg J. Flesch¹ and Ken A. Farley⁴



¹Jet Propulsion Laboratory, California Institute of Technology, Pasadena CA 91109






²NASA Goddard Space Flight Center (GSFC), Greenbelt, MA 20771

³University of Michigan, Ann Arbor, MI 48109



⁴California Institute of Technology, Pasadena CA 91125

The Tunable Laser Spectrometer (TLS) is one of three instruments that make up the Sample Analysis at Mars (SAM) suite on the Curiosity Rover that landed in August 2012. TLS is a two-channel tunable laser spectrometer (3.7 kg) using an Interband Cascade (IC) laser at 3.27 μm for methane measurements, and a near-IR tunable diode laser for measurements of water and carbon dioxide isotopes. To date, TLS has measured in CO_2 the isotope ratios $^{13}\text{C}/^{12}\text{C}$, $^{18}\text{O}/^{16}\text{O}$, $^{17}\text{O}/^{16}\text{O}$ and $^{13}\text{C}^{18}\text{O}/^{12}\text{C}^{16}\text{O}$; and in water the isotope ratios D/H and $^{18}\text{O}/^{16}\text{O}$ in both the atmosphere and gases evolved from pyrolysis of soils and rock samples. Only methane search results are reported here.

Methane in the atmosphere of Mars is a potential signature of ongoing or past biological activity on the planet. During the last decade, Earth-based telescopic and Mars orbit remote sensing instruments have reported significant abundances of methane in the Martian atmosphere ranging from several to tens of parts-per-billion by volume (ppbv). Observations from Earth showed "plumes" of methane with variations on timescales much faster than expected and inconsistent with localized patches seen from orbit, prompting speculation of sources from sub-surface methanogen bacteria, geological water-rock reactions or infall from comets, micro-meteorites or interplanetary dust. From measurements on NASA's Curiosity Rover that landed near Gale Crater on 5th August 2012, we here report **no definitive detection of methane** in the near-surface Martian atmosphere. Our *in situ* measurements were made using the Tunable Laser Spectrometer (TLS) in the Sample Analysis at Mars (SAM) instrument suite that made three separate searches on Martian sols 79, 81 and 106 after landing. The measured mean value of 0.4 ppbv corresponds to an **upper limit for methane abundance of <3 ppbv** at the 95% confidence level.



The IC laser was invented (Rui Yang et al.) and developed by researchers at JPL's Microdevices Laboratory (MDL). For TLS, it produces single-mode output of ~5 mW at 245 K. The laser is scanned over the methane region every second with spectra co-averaged for 2 mins on board before downloading to Earth. The TLS NIR laser was provided by Nanoplus.



Mars methane at Gale Crater (4.5°S, 137°E, Ls=157): Expectations

Technique	Observation	Max (ppbv)	Min (ppbv)	Prediction (ppbv)
TLS-SAM	v3/R3 (3057.7 cm^{-1})	3 ppbv (2 σ)	-	< 3 ppbv
MEX/PFS	v3/Q (3018 cm^{-1})	70	< 3	10-15 ppbv
2003- Present	R=2000	N. summer	N. winter	
MGS/TES	v3/Q (1306 cm^{-1})	N. polar 60	< 1	5-20
2001-2003	R=100-200	N. summer	N. winter	
IRTf, Keck	v3/(R0, R1 at 3029 and 3039 cm^{-1})	N. polar 40	< 3	20-30
2003-2006	R=10 ⁵	N. summer Equatorial	N. winter	
CFHT/FTS	v3/P (2997 cm^{-1})	10 (<30)	10 (<30)	10 (<30)
2003	R=1.8X10 ⁵	Global average	Global average	Global average

REFERENCES

Atreya, S. K. et al. Methane on Mars: Current observations, interpretation, and future plans. *Planet. Space Sci.* **59**, 133-136 (2011).
 Atreya, S. K. et al. Strong release of methane on Mars in northern summer 2003. *Science* **323**, 1041-1045 (2009).
 Atreya, S. K., Encrenaz, T., Ignatiev, N. & Garavito, M. Detection of methane in the atmosphere of Mars. *Nature* **386**, 1758-1761 (2004).
 Commins, A., Fortmann, V. & Sandoz, G. Mapping methane in Martian atmosphere with PFS-MEX data. *Planet. Space Sci.* **99**, 172-148 (2011).
 Mahaffy, P. R. et al. The Sample Analysis at Mars Investigation and Instrument Suite. *Space Sci. Rev.* **170**, doi:10.1007/11214-012-9879-9 (2012).
 Patel, S. & Martin, G. K. Mapping the methane on Mars. *Atmos. Environ.* **42**, 451-461 (2008).
 Villanueva, G. L. et al. A sensitive search for Organics (CH₄, CH₃OH, H₂CO, C₂H₆, C₂H₄, hydroperoxides/RO₂), inorganic compounds (N₂O, NH₃, HCN) and chlorine species (CH₃Cl, CH₂Cl) on Mars using ground-based high-resolution infrared spectroscopy. *Atmos. Environ.* **22A**, 11-27 (2011).
 Atreya, S. K. et al. Search for methane and upper limits to ethane and SO₂ on Mars. *Astron. J.* **137**, 144-151 (2012).
 Atreya, S. K. & Forget, F. Observed variations of methane on Mars unexplained by known atmospheric chemistry and physics. *Nature* **440**, doi:10.1038/nature08228 (2009).
 Zaleski, K. J., Fortmann, V. & Sandoz, G. C. Is there methane on Mars? *Planet. Space Sci.* **112**, 493-501 (2011).
 Webster, C. R. & Mahaffy, P. R. Determining the local abundance of Martian methane and its ¹³C/¹²C and D/H isotopic ratios for comparison with related gas and soil analysis in the 2013 Mars Science Laboratory (MSL) mission. *Planet. Space Sci.* **99**, 271-280 (2011).
 Wong, A. S., Atreya, S. K. & Encrenaz, T. Chemical markers of possible hot spots on Mars. *J. Geophys. Res.* **118**, doi:10.1029/2002JE002003 (2003).
 Atreya, S. K. et al. Oxidant Enhancement in Martian Dust Devils and Storms: Implications for Life, and Habitability. *Astrobiology* **4**, 479-501 (2004).
 Chéry, G. et al. Oxidant Enhancement in Martian Dust Devils and Storms: Storm Electric Fields and Electron Discharge. *Astrobiology* **4**, 451-462 (2004).
 Farrell, W. M., Delory, G. T., Atreya, S. K. Martian Dust Storms as a Possible Sink of Atmospheric Methane. *J. Geophys. Res.* **33**, doi:10.1029/2006GL027710 (2006).
 Atreya, S. K., Mahaffy, P. R. & Wong, A. S. Methane & related trace species on Mars: Origin, loss, implications for life, & habitability. *Planet. Space Sci.* **85**, 538-569 (2007).
 Kassel, E. Measurable methane sulfate particles as a source of methane to the martian atmosphere. *Icarus* **204**, 137-144 (2009).
 Atreya, S. K. Ultraviolet radiation-induced methane emissions from meteorites and the Martian atmosphere. *Nature* **482**, 95-96, doi:10.1038/nature11201 (2012).
 Schaefer, A. C., Moons, J. E., Claux, C. A., Barlow, N. G. & Britt, D. T. Methane from UV-irradiated carbonaceous chondrites under simulated Martian conditions. *J. Geophys. Res.* **117**, doi:10.1029/2011JB004125 (2012).

Acknowledgement: The research described here was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration (NASA). The TLS instrument was built at JPL and delivered to NASA GSFC for integration into the SAM suite.

