

**EXTREME H ISOTOPIC ANOMALIES IN CHONDRITIC ORGANIC MATTER.** H. Busemann<sup>1</sup>, C. M. O'D. Alexander<sup>1</sup>, P. Hoppe<sup>2</sup>, L. R. Nittler<sup>1</sup>, and A. F. Young<sup>1</sup>. <sup>1</sup>DTM, Carnegie Institution of Washington, Washington DC, 20015, USA. E-mail: busemann@dtm.ciw.edu. <sup>2</sup>MPI for Chemistry, Mainz, Germany.

**Introduction:** Organic matter in meteorites and interplanetary dust particles (IDPs) show isotope enrichments in D and <sup>15</sup>N testifying to the partial origin of this material in the protosolar cloud [1]. Spatial correlations, carriers and associated minerals of these anomalies are not well characterized. We have conducted ionprobe (IMS6f and NanoSIMS) imaging studies of various samples for H, D, C, <sup>14</sup>N and <sup>15</sup>N. These will ultimately be correlated with micro-analytic techniques such as FIB/TEM or STXM/XANES. We analyzed matrix fragments from Bells (CM2), Al Rais (CR2) and Tagish Lake (unique) [2], high purity insoluble organic matter (IOM) [3] extracted from EET92042 ("EET", CR2), Bells, Murchison (CM2), Allende (CV3), Krymka (LL3.1) and, for comparison, 3 IDPs.

We have found extreme enrichments in D in the IOM of EET, <sup>15</sup>N excesses (correlated and uncorrelated with D) in the same material and in Bells matrix fragments [4]. These anomalies exceed all but one of the enriched hotspots found in IDPs. Hence, the IOM in primitive meteorites retained a memory of interstellar chemistry that is comparable to that found in IDPs.

**Ionprobe calibration:** We measured a range of well-characterized organics including coals, a lipid and meteoritic IOM [3] to assess potential matrix effects that may hamper the *absolute* determination of D/H and C/H by ionprobe raster imaging with the IMS6f ionprobe. While the instrumental mass fractionation for D/H is within 15 % for all samples, we found variations in C/H that depend on the samples' elemental compositions. Samples with low C/H (e.g. EET) are well behaved, but those with large C/H (e.g. Allende) have proved difficult to calibrate.

**Results and Discussion:** While the bulk IOM in EET is  $\delta D \sim 3000$  ‰ [3], about 1 area% of the material exhibits larger D isotope anomalies. We found 22 regions (diameter  $\geq 1.4$   $\mu m$ ,  $\sim$ beam size) that show  $\delta D$  values of 5000–16000 ‰. C/H in these regions is comparable to the bulk IOM value. D "hotspots" were also found in the Murchison IOM, reaching values of  $\sim 1700$  ‰, indicating that some primitive material has survived the parent-body processes that have modified the IOM in CM chondrites [3]. The highest  $\delta D$  values ( $\sim 16000$  ‰) in the EET IOM exceed those found in all but one IDP [5,6] and approach the  $\delta D$  value obtained for cometary HCN [7]. This IOM is an agglomerate of matter assembled by the demineralization of bulk rock. The actual D "hotspots" could well be much smaller than our beam size and the  $\delta D$  accordingly higher. The results show that the parent bodies of the most primitive carbonaceous meteorites acquired an assemblage of IOM as primitive as the parent bodies of the IDPs.

**References:** [1] Ehrenfreund P. & Charnley S. B. 2000 *Annual Review Astronomy Astrophysics* 38:427-483. [2] Young A. F. et al. 2004. #2097. 35th Lunar Planetary Science Conference. [3] Alexander C. M. O'D. et al. 2005. *Meteoritics and Planetary Science* 40: this volume. [4] Nittler L.R. et al. 2005. *Meteoritics and Planetary Science* 40: this volume. [5] Aléon J. et al. 2001. *Geochimica and Cosmochimica Acta* 65: 4399-4412. [6] Messenger S. 2000. *Nature* 404: 968-971. [7] Meier R. et al. 1998 *Science* 279: 1707-1710.